

## BLACKBODY RADIATION: CLASSICAL APPROXIMATIONS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

Reference: Carroll, Bradley W. & Ostlie, Dale A. (2007), *An Introduction to Modern Astrophysics*, 2nd Edition; Pearson Education - Chapter 3, Problems 3.10 - 3.12.

The blackbody radiation rate in terms of wavelength is

$$(1) \quad B_{\lambda}(T) = \frac{2hc^2}{\lambda^5 (e^{hc/\lambda k_B T} - 1)}$$

For very short wavelengths (high energies),  $e^{hc/\lambda k_B T} \gg 1$  so we can approximate the rate by

$$(2) \quad B_{\lambda} \approx \frac{2hc^2}{\lambda^5} e^{-hc/\lambda k_B T}$$

This form matches the empirical formula obtained by Wien, which is

$$(3) \quad B_{\lambda} = \frac{a}{\lambda^5} e^{-b/T}$$

where  $a$  and  $b$  are constants determined by fitting the curve to experimental data.


For long wavelengths, we can use the first order Taylor expansion for the exponential

$$(4) \quad e^{hc/\lambda k_B T} = 1 + \frac{hc}{\lambda k_B T} + \mathcal{O}\left(\frac{1}{\lambda^2}\right)$$

$$(5) \quad B_{\lambda} \approx \frac{2ck_B T}{\lambda^4}$$

This is the classical Rayleigh-Jeans law. Notice that Planck's constant has dropped out of the formula, and that if applied to all wavelengths it predicts that  $B_{\lambda} \rightarrow \infty$  as  $\lambda \rightarrow 0$ . This is the *ultraviolet catastrophe* which was rectified with the introduction of the quantization of energy.

To compare the three curves for the Sun, where  $T = 5777$  K, we have the following plot:



The green curve is Planck's formula, the blue curve is the Rayleigh-Jeans formula and the red curve is Wien's empirical formula. The Rayleigh-Jeans formula predicts twice the actual radiation at around  $\lambda = 2000$  nm which is in the infrared region.

We've already derived Wien's displacement law for the wavelength at which the blackbody curve is maximum:

$$(6) \quad \lambda_{max} = \frac{2.901 \times 10^{-3}}{T} \text{ m}$$