

## RELATIVISTIC ACCELERATION IN TERMS OF FORCE

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: Carroll, Bradley W. & Ostlie, Dale A. (2007), *An Introduction to Modern Astrophysics*, 2nd Edition; Pearson Education - Chapter 4, Problem 4.14.

Ordinary force in relativity is given by

$$(0.1) \quad \mathbf{F} = \frac{m}{\sqrt{1-u^2/c^2}} \left[ \mathbf{a} + \frac{(\mathbf{u} \cdot \mathbf{a}) \mathbf{u}}{c^2 - u^2} \right]$$

To get a general expression for the acceleration  $\mathbf{a}$  in terms of the force, we take the dot product of both sides with the velocity  $\mathbf{u}$ :

$$(0.2) \quad \mathbf{u} \cdot \mathbf{F} = \gamma m (\mathbf{u} \cdot \mathbf{a}) \left( 1 + \frac{u^2}{c^2 - u^2} \right)$$

$$(0.3) \quad = \frac{\gamma m c^2 (\mathbf{u} \cdot \mathbf{a})}{c^2 - u^2}$$

$$(0.4) \quad = \gamma^3 m (\mathbf{u} \cdot \mathbf{a})$$

$$(0.5) \quad \mathbf{u} \cdot \mathbf{a} = \frac{\mathbf{u} \cdot \mathbf{F}}{\gamma^3 m}$$

Substituting back into 0.1, we get

$$(0.6) \quad \mathbf{a} = \frac{\mathbf{F}}{\gamma m} - \frac{\mathbf{u}}{c^2 - u^2} \frac{\mathbf{u} \cdot \mathbf{F}}{\gamma^3 m}$$

$$(0.7) \quad = \frac{\mathbf{F}}{\gamma m} - \frac{\mathbf{u}}{\gamma m c^2} (\mathbf{u} \cdot \mathbf{F})$$

In the limit of small  $\mathbf{u}$ , this reduces to the familiar Newton's law  $\mathbf{F} = m\mathbf{a}$ , but in the relativistic region, the acceleration depends on the object's velocity. As a result, the acceleration isn't parallel to the force unless  $\mathbf{F}$  is either parallel to  $\mathbf{u}$  or  $\mathbf{F} \perp \mathbf{u}$ ; in the latter case  $\mathbf{u} \cdot \mathbf{F} = 0$  and the second term is zero.

PINGBACKS

Pingback: Acceleration under a constant force