

SIRIUS BINARY STAR SYSTEM

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

Reference: Carroll, Bradley W. & Ostlie, Dale A. (2007), *An Introduction to Modern Astrophysics*, 2nd Edition; Pearson Education - Chapter 7, Problem 7.4.

As an example of using measurements of a binary star system's properties to determine some of its physical characteristics, we'll look at Sirius. Sirius is a visual binary (that is, both stars are visible directly) with a period of $P = 49.94$ years. The parallax of Sirius is $p = 0.37921'' \pm 0.00158''$. If we take the inclination angle to be $i = 0^\circ$, that is, the orbital plane of the Sirius system is in the plane of the sky, then the angular semimajor axis of the reduced mass is $\alpha = 7.61''$. The ratio of the distances of Sirius A (the bright star) and Sirius B (the white dwarf companion) from the centre of mass is $a_A/a_B = 0.466$.

From the parallax, we can get the distance:

$$(0.1) \quad d = \frac{1}{p} = 2.637 \text{ pc} = 8.597 \text{ ly}$$

We can then find the masses of the two stars. The mass ratio is

$$(0.2) \quad \frac{m_B}{m_A} = \frac{a_A}{a_B} = 0.466$$

From Kepler's third law we can get the sum of the masses:

$$(0.3) \quad m_A + m_B = \frac{4\pi^2}{GP^2} a^3$$

where a is the semimajor axis of the centre of mass. Since we know the angular semimajor axis and the distance, we have

$$(0.4) \quad a = \alpha d$$

$$(0.5) \quad = 7.61'' \frac{\pi}{3600 \times 180} \times 8.597$$

$$(0.6) \quad = 3.172 \times 10^{-4} \text{ ly}$$

$$(0.7) \quad = 3.00 \times 10^{12} \text{ m}$$

The period is $P = 1.576 \times 10^9$ s so the sum of the masses is

$$(0.8) \quad m_A + m_B = \frac{4\pi^2 (3.00 \times 10^{12})^3}{(6.67 \times 10^{-11})(1.576 \times 10^9)^2}$$

$$(0.9) \quad = 6.434 \times 10^{30} \text{ kg}$$

$$(0.10) \quad = 3.235M_S$$

where M_S is the mass of the Sun. From this and 0.2, we get

$$(0.11) \quad m_B = 0.466m_A$$

$$(0.12) \quad m_A = \frac{3.235}{1.466}M_S$$

$$(0.13) \quad = 2.21M_S$$

$$(0.14) \quad m_B = 1.03M_S$$

[These values are somewhat higher than the currently accepted values of $m_A = 2.02M_S$ and $m_B = 0.978M_S$. The discrepancy arises from the currently measured angular semimajor axis being smaller at $\alpha = 7.50 \pm 0.04''$, the period is slightly longer at $P = 50.09 \pm 0.055$ yr and the angle of inclination not being 0, but rather $i = 136.53 \pm 0.43^\circ$.]

We've already worked out the luminosity of Sirius A from its absolute bolometric magnitude of $M_A = +1.36$ and that of the Sun ($M_S = +4.74$):

$$(0.15) \quad \frac{L_A}{L_S} = 100^{(M_S - M_A)/5} = 22.49$$

For the companion $M_B = +8.79$:

$$(0.16) \quad \frac{L_B}{L_S} = 100^{(M_S - M_B)/5} = 0.024$$

Given the surface temperature of Sirius B as $T_B = 24,790$ K and treating it as a blackbody, we can estimate its radius R_B from the formula

$$(0.17) \quad L_B = 4\pi R_B^2 \sigma T_B^4$$

where $\sigma = 5.670373 \times 10^{-8} \text{ W m}^{-2}\text{K}^{-4}$ is the Stefan-Boltzmann constant. The luminosity of the Sun is

$$(0.18) \quad L_S = 3.839 \times 10^{26} \text{ watts}$$

so

$$(0.19) \quad R_B = \sqrt{\frac{L_B}{4\pi\sigma T_B^4}}$$

$$(0.20) \quad = \sqrt{\frac{0.024L_S}{4\pi\sigma T_B^4}}$$

$$(0.21) \quad = \sqrt{\frac{0.024 \times 3.839 \times 10^{26}}{4\pi(5.670373 \times 10^{-8})(24790)^4}}$$

$$(0.22) \quad = 5.85 \times 10^6 \text{ m}$$

This is slightly less than the radius of the Earth (6.371×10^6 m) and only 0.0084 times the radius of the Sun (6.963×10^8 m).