

MAXWELL'S EQUATIONS IN TERMS OF POTENTIALS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog and include the title or URL of this post in your comment.

Post date: 3 August 2021.

Maxwell's equations are

$$\nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0} \quad (1)$$

$$\nabla \cdot \mathbf{B} = 0 \quad (2)$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \quad (3)$$

$$\nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} \quad (4)$$

They are completely general, in that they describe the interactions between electric and magnetic fields and the charges and currents that generate them, in both the static (fixed charge and current densities) and dynamic (time-varying) cases. In the static case, the two time derivatives are zero, which means that $\nabla \times \mathbf{E} = 0$. Since any vector field whose curl is zero can be represented as the gradient of a scalar field, we could write $\mathbf{E} = -\nabla V$ where V is the potential. The presence of the term $-\frac{\partial \mathbf{B}}{\partial t}$, however, means that we can't, in general, write \mathbf{E} as the gradient of a scalar field in the dynamic case.

Because $\nabla \cdot \mathbf{B} = 0$ we can write \mathbf{B} as the curl of a vector field

$$\mathbf{B} = \nabla \times \mathbf{A} \quad (5)$$

This is still true in the dynamic case. Substituting this into 3 we get

$$\nabla \times \mathbf{E} = -\frac{\partial \nabla \times \mathbf{A}}{\partial t} \quad (6)$$

$$\nabla \times \left(\mathbf{E} + \frac{\partial \mathbf{A}}{\partial t} \right) = 0 \quad (7)$$

The LHS is the curl of a vector field that combines the electric field and the magnetic vector potential, and since it is zero even in the dynamic case, we can write this vector field as the gradient of a scalar field V . [We should be able to use the same symbol V for the potential in the dynamic case,

since if $\frac{\partial \mathbf{A}}{\partial t} = 0$ we regain the static equations $\nabla \times \mathbf{E} = 0$ and $\mathbf{E} = -\nabla V$.] That is, we can write

$$\mathbf{E} + \frac{\partial \mathbf{A}}{\partial t} = -\nabla V \quad (8)$$

$$\mathbf{E} = -\nabla V - \frac{\partial \mathbf{A}}{\partial t} \quad (9)$$

Equations 5 and 9 effectively replace 2 and 3. Using 5 and 9 we can eliminate \mathbf{E} and \mathbf{B} from 1 and 4:

$$\nabla \cdot \mathbf{E} = -\nabla^2 V - \frac{\partial}{\partial t} (\nabla \cdot \mathbf{A}) \quad (10)$$

$$\nabla^2 V + \frac{\partial}{\partial t} (\nabla \cdot \mathbf{A}) = -\frac{\rho}{\epsilon_0} \quad (11)$$

$$\nabla \times \mathbf{B} = \nabla \times (\nabla \times \mathbf{A}) \quad (12)$$

$$= \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} \quad (13)$$

$$\nabla \times (\nabla \times \mathbf{A}) = \mu_0 \mathbf{J} - \mu_0 \epsilon_0 \nabla \frac{\partial V}{\partial t} - \mu_0 \epsilon_0 \frac{\partial^2 \mathbf{A}}{\partial t^2} \quad (14)$$

We can rearrange the last equation using the identity

$$\nabla \times (\nabla \times \mathbf{A}) = \nabla (\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A} \quad (15)$$

so we get

$$\nabla (\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A} = \mu_0 \mathbf{J} - \mu_0 \epsilon_0 \nabla \frac{\partial V}{\partial t} - \mu_0 \epsilon_0 \frac{\partial^2 \mathbf{A}}{\partial t^2} \quad (16)$$

$$\nabla^2 \mathbf{A} - \mu_0 \epsilon_0 \frac{\partial^2 \mathbf{A}}{\partial t^2} - \nabla \left(\nabla \cdot \mathbf{A} + \mu_0 \epsilon_0 \frac{\partial V}{\partial t} \right) = -\mu_0 \mathbf{J} \quad (17)$$

To summarize:

$$\boxed{\nabla^2 V + \frac{\partial}{\partial t} (\nabla \cdot \mathbf{A}) = -\frac{\rho}{\epsilon_0}} \quad (18)$$

$$\boxed{\nabla^2 \mathbf{A} - \mu_0 \epsilon_0 \frac{\partial^2 \mathbf{A}}{\partial t^2} - \nabla \left(\nabla \cdot \mathbf{A} + \mu_0 \epsilon_0 \frac{\partial V}{\partial t} \right) = -\mu_0 \mathbf{J}} \quad (19)$$

These two equations comprise 4 equations (one from 18 and one for each vector component in 19) for four functions (V and \mathbf{A}), and their solution allows us to calculate both \mathbf{E} and \mathbf{B} by means of 9 and 5, so they form a

complete replacement for the original set of 4 Maxwell equations that we started with. This reduces the number of equations to be solved from 6 (for the 3 components of each of \mathbf{E} and \mathbf{B}) to 4.

We can write these equations in a more compact form using the d'Alembertian operator

$$\square^2 \equiv \nabla^2 - \mu_0 \epsilon_0 \frac{\partial^2}{\partial t^2} \quad (20)$$

and defining the function

$$L \equiv \nabla \cdot \mathbf{A} + \mu_0 \epsilon_0 \frac{\partial V}{\partial t} \quad (21)$$

We get

$$\square^2 V = \nabla^2 V - \mu_0 \epsilon_0 \frac{\partial^2 V}{\partial t^2} \quad (22)$$

$$= \nabla^2 V - \frac{\partial L}{\partial t} + \frac{\partial}{\partial t} (\nabla \cdot \mathbf{A}) \quad (23)$$

$$\square^2 V + \frac{\partial L}{\partial t} = \nabla^2 V + \frac{\partial}{\partial t} (\nabla \cdot \mathbf{A}) \quad (24)$$

$$= -\frac{\rho}{\epsilon_0} \quad (25)$$

using 18 in the last line. Also,

$$\square^2 \mathbf{A} = \nabla^2 \mathbf{A} - \mu_0 \epsilon_0 \frac{\partial^2 \mathbf{A}}{\partial t^2} \quad (26)$$

$$\square^2 \mathbf{A} - \nabla L = -\mu_0 \mathbf{J} \quad (27)$$

using 19. In summary

$$\boxed{\square^2 V + \frac{\partial L}{\partial t} = -\frac{\rho}{\epsilon_0}} \quad (28)$$

$$\boxed{\square^2 \mathbf{A} - \nabla L = -\mu_0 \mathbf{J}} \quad (29)$$

PINGBACKS

Pingback: Energy flow in time-dependent fields

Pingback: Potentials for a point charge

Pingback: Potentials for an electromagnetic wave

Pingback: Retarded potentials in an infinite straight wire

Pingback: Jefimenko's equation for time-dependent electric field

Pingback: Fields of a moving point charge

Pingback: Electric dipole time-varying potentials - Lorenz gauge condition

Pingback: Lagrangian for classical electromagnetism

Pingback: Euler-Lagrange equations for a free particle in an electromagnetic field

Pingback: Maxwell's equations in 4d spacetime form

Pingback: Lorenz gauge in 4d spacetime form

Pingback: Maxwell's equations with sources in 4d spacetime