

POTENTIALS FOR A POINT CHARGE

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog and include the title or URL of this post in your comment.

Post date: 3 August 2021.

As a simple (though unusual) example of specifying a system through electric and magnetic potentials suppose we have

$$V(\mathbf{r}, t) = 0 \quad (1)$$

$$\mathbf{A}(\mathbf{r}, t) = -\frac{1}{4\pi\epsilon_0} \frac{qt}{r^2} \hat{\mathbf{r}} \quad (2)$$

These potentials give rise to the fields

$$\mathbf{B} = \nabla \times \mathbf{A} = 0 \quad (3)$$

$$\mathbf{E} = -\nabla V - \frac{\partial \mathbf{A}}{\partial t} = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{\mathbf{r}} \quad (4)$$

The expression for \mathbf{E} is just that of a point charge q at the origin, while a zero magnetic field indicates that there is no current. If we want to be pedantic, we can also get these results from the potentials. For the charge density, we had

$$\nabla^2 V + \frac{\partial}{\partial t} (\nabla \cdot \mathbf{A}) = -\frac{\rho}{\epsilon_0} \quad (5)$$

Using the formula

$$\nabla \cdot \left(\frac{1}{r^2} \hat{\mathbf{r}} \right) = 4\pi \delta_3(\mathbf{r}) \quad (6)$$

we get

$$\nabla \cdot \mathbf{A} = -\frac{qt}{\epsilon_0} \delta_3(\mathbf{r}) \quad (7)$$

$$\rho = q\delta_3(\mathbf{r}) \quad (8)$$

For the current, we can use the equation

$$\nabla \times (\nabla \times \mathbf{A}) = \mu_0 \mathbf{J} - \mu_0 \epsilon_0 \nabla \frac{\partial V}{\partial t} - \mu_0 \epsilon_0 \frac{\partial^2 \mathbf{A}}{\partial t^2} \quad (9)$$

and we see that all the terms not involving \mathbf{J} are zero, so $\mathbf{J} = 0$ as well. Thus this is a bizarre way of writing potentials for a point charge. This illustrates that the potentials giving rise to a particular charge and current distribution are not unique.

PINGBACKS

Pingback: [Gauge transformations in electrodynamics](#)

Pingback: [Coulomb and Lorenz gauges](#)