

## LORENTZ TRANSFORMATIONS AND THE SPECIAL LINEAR GROUP $SL(2,\mathbb{C})$

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: W. Greiner & J. Reinhardt, *Field Quantization*, Springer-Verlag (1996), Chapter 2, Section 2.4.

Arthur Jaffe, *Lorentz transformations, rotations and boosts*, online notes available (at time of writing, Sep 2016) here.

Continuing our examination of general Lorentz transformations, we start off with the representation of a spacetime 4-vector as a  $2 \times 2$  complex Hermitian matrix:

$$\hat{x} \equiv \begin{bmatrix} x_0 + x_3 & x_1 - ix_2 \\ x_1 + ix_2 & x_0 - x_3 \end{bmatrix} \quad (1)$$

Our ultimate goal is to show that any Lorentz transformation can be represented as the product of a pure rotation  $R$  and a pure boost  $B$ :  $\Lambda = RB$ . The step shown in this post may look like little more than an exercise in matrix algebra, but be patient; it takes a while to get to our final goal.

We start by looking at the matrices belonging to the special linear group  $SL(2, \mathbb{C})$ , which consists of  $2 \times 2$  matrices containing general complex numbers as elements, and with determinant 1. Each matrix  $A \in SL(2, \mathbb{C})$  can be used to define a linear transformation of the Hermitian matrix 1:

$$\hat{x}' = A\hat{x}A^\dagger \quad (2)$$

Because the determinant of a product is equal to the product of the determinants, and  $\det A = \det A^\dagger = 1$ ,  $\det \hat{x}' = \det \hat{x} = x_\mu x^\mu$ . Thus such a transformation leaves the 4-vector length unchanged, so qualifies as a Lorentz transformation. Also, as a general complex  $2 \times 2$  matrix contains 4 elements, each with a real and imaginary part, there are 8 parameters. The condition  $\det A = 1$  provides 2 constraints (one on the real part and one on the imaginary part), leaving 6 independent parameters, which is the same as the number of free parameters in a general Lorentz transformation.

We can give a more detailed proof that  $A$  provides a Lorentz transformation as follows. Suppose we start with two matrices  $A, B \in SL(2, \mathbb{C})$  and define a transformation

$$\hat{x}' = A\hat{x}B \quad (3)$$

[Remember that the hats on  $\hat{x}$  and  $\hat{x}'$  mean that we're considering the  $2 \times 2$  matrix version 1 of the 4-vectors  $x$  and  $x'$ .] The transformed matrix  $\hat{x}'$  must be Hermitian for all  $\hat{x}$ , so we must have

$$(A\hat{x}B)^\dagger = A\hat{x}B \quad (4)$$

$$= B^\dagger \hat{x} A^\dagger \quad (5)$$

We now left-multiply by  $(B^\dagger)^{-1}$  and right-multiply by  $B^{-1}$  to get

$$(B^\dagger)^{-1} A \hat{x} = \hat{x} A^\dagger B^{-1} \quad (6)$$

But we also have

$$(B^\dagger)^{-1} A = (A^\dagger B^{-1})^\dagger \quad (7)$$

so the matrix

$$T \equiv (B^\dagger)^{-1} A \quad (8)$$

is Hermitian. We can therefore write 6 as

$$T \hat{x} = \hat{x} T^\dagger = \hat{x} T \quad (9)$$

so  $T$  commutes with  $\hat{x}$  for all  $\hat{x}$ .

Now we can choose  $x = \sigma_2$  and then  $x = \sigma_3$ , where the  $\sigma_i$ s are two of the Pauli matrices which we showed (together with the identity matrix  $\sigma_0$ ) form a basis for the space of  $2 \times 2$  Hermitian matrices. Now we've seen that  $\sigma_2$  and  $\sigma_3$  also form an irreducible set, and we saw that any matrix  $T$  that commutes with all the members of an irreducible set must be a multiple of the identity matrix. Thus we must have

$$T = \lambda I \quad (10)$$

for some constant  $\lambda$ . However, since  $T$  is the product of two matrices  $A$  and  $(B^\dagger)^{-1}$ , both of which have determinant 1,  $\det T = 1$  also, which means that  $\lambda^2 = 1$  and  $\lambda = \pm 1$ . Therefore

$$(B^\dagger)^{-1} A = \pm I \quad (11)$$

$$A = \pm B^\dagger \quad (12)$$

Thus the transformation 3 can be written as

$$\hat{x}' = \pm A \hat{x} A^\dagger \quad (13)$$

To eliminate the  $-$  sign, suppose that

$$\widehat{x}' = -A\widehat{x}A^\dagger \quad (14)$$

A Lorentz transformation giving this result can be written as

$$\widehat{x}' = \widehat{\Lambda x} \quad (15)$$

where  $\Lambda$  is the  $4 \times 4$  matrix giving the Lorentz transformation of the original 4-vector  $x$ . In the original 4-vector notation, we have

$$x'_\mu = \sum_{\nu=0}^3 \Lambda_{\mu\nu} x_\nu \quad (16)$$

$$= (\Lambda x)_\mu \quad (17)$$

From the relation between the 4-vector and  $2 \times 2$  matrix representations, we have

$$x'_\mu = \langle \sigma_\mu, \widehat{x}' \rangle \quad (18)$$

where  $\langle \sigma_\mu, \widehat{x}' \rangle$  is the inner product of the two matrices. Therefore from 14

$$(\Lambda x)_\mu = \langle \sigma_\mu, \widehat{x}' \rangle \quad (19)$$

$$= -\langle \sigma_\mu, A\widehat{x}A^\dagger \rangle \quad (20)$$

$$= -\left\langle \sigma_\mu, A \left( \sum_{\nu=0}^3 \sigma_\nu x_\nu \right) A^\dagger \right\rangle \quad (21)$$

If we choose  $x = (1, 0, 0, 0)$ , we have

$$(\Lambda x)_0 = \Lambda_{00} \quad (22)$$

$$= -\left\langle \sigma_0, A \left( \sum_{\nu=0}^3 \sigma_\nu x_\nu \right) A^\dagger \right\rangle \quad (23)$$

$$= -\langle \sigma_0, A\sigma_0A^\dagger \rangle \quad (24)$$

$$= -\langle \sigma_0, AA^\dagger \rangle \quad (25)$$

$$= -\frac{1}{2} \text{Tr} (AA^\dagger) \quad (26)$$

$$\leq 0 \quad (27)$$

where the penultimate line follows from the definition of the inner product. The last line follows because

$$\text{Tr}(AA^\dagger) = |A_{11}|^2 + |A_{22}|^2 \geq 0 \quad (28)$$

Since we're requiring the transformation to be orthochronous, we must have  $\Lambda_{00} \geq 1$ , so we must exclude the  $-$  sign in 13, giving 2.

Finally, we can show that the transformation matrix  $A$  is unique, up to a sign. We can prove this by supposing that there are two different  $SL(2, \mathbb{C})$  matrices  $A$  and  $B$  that give the same transformation for all  $\hat{x}$ , that is

$$A\hat{x}A^\dagger = B\hat{x}B^\dagger \quad (29)$$

This implies

$$B^{-1}A\hat{x}A^\dagger(B^\dagger)^{-1} = \hat{x} \quad (30)$$

$$= B^{-1}A\hat{x}(B^{-1}A)^\dagger \quad (31)$$

We can now choose  $\hat{x} = I$ , which shows that

$$(B^{-1}A)^\dagger = (B^{-1}A)^{-1} \quad (32)$$

which means (by definition),  $B^{-1}A$  is unitary, so for all  $\hat{x}$

$$\hat{x} = B^{-1}A\hat{x}(B^{-1}A)^{-1} \quad (33)$$

This means that  $B^{-1}A$  commutes with  $\hat{x}$  for all  $\hat{x}$  (that's the only way we can cancel  $B^{-1}A$  off the RHS). Using the same argument as above, we can choose  $\hat{x}$  to be two of the Pauli matrices, which form an irreducible set. Since  $B^{-1}A$  commutes with both these matrices, it must be a multiple  $\lambda$  of the identity:

$$B^{-1}A = \lambda I \quad (34)$$

$$A = \lambda B \quad (35)$$

Since  $\det A = \det B = 1$  and for a  $2 \times 2$  matrix  $\det(\lambda B) = \lambda^2 \det B$ , we have  $\lambda^2 = 1$ , so  $\lambda = \pm 1$ . Therefore  $A$  is unique up to a sign.

In summary, what we've done in this post is show that a restricted Lorentz transformation  $\Lambda$  (that is, one where  $\det \Lambda = +1$  and  $\Lambda_{00} \geq 1$ ) can be represented by a matrix  $A \in SL(2, \mathbb{C})$  where  $A$  is unique up to a sign.

PINGBACKS

Pingback: Lorentz transformation as product of a pure boost and pure rotation