

KLEIN-GORDON EQUATION IN SCHRÖDINGER FORM - LAGRANGIAN, ENERGY-MOMENTUM TENSOR

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: W. Greiner: *Relativistic Quantum Mechanics (Wave Equations)*; 3rd Edition, Springer-Verlag (2000); Section 1.8; Exercise 1.7.

Post date: 10 Dec 2017.

Greiner's Exercise 1.7 introduces the Lagrangian density and energy-momentum tensor for the Schrödinger representation of the Klein-Gordon equation. (In the text, he says that it's for the Feshbach-Villars representation, but all the equations refer to the two-component wave function Ψ rather than the function Φ which is used in the Feshbach-Villars form.)

As usual, the Lagrangian density is just stated (one of these days I'd like to find out if it is possible to actually *derive* a Lagrangian rather than conjuring them out of thin air) to be

$$\mathcal{L} = i\hbar\bar{\Psi}\partial_t\Psi - \frac{\hbar^2}{2m_0}\nabla\bar{\Psi}(\tau_3 + i\tau_2)\nabla\Psi - m_0c^2\bar{\Psi}\tau_3\Psi \quad (1)$$

where the τ_i matrices are essentially the Pauli matrices

$$\tau_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}; \quad \tau_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}; \quad \tau_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad \mathbf{1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad (2)$$

and

$$\bar{\Psi} \equiv \Psi^\dagger \tau_3 \quad (3)$$

By using the usual technique of varying the action integral, Greiner shows that the Euler-Lagrange equations yield the Schrödinger representation of the Klein-Gordon equation. That is, variation of $\bar{\Psi}$ yields the equation

$$\partial_\nu \frac{\partial \mathcal{L}}{\partial (\partial_\nu \bar{\Psi}_\alpha)} - \frac{\partial \mathcal{L}}{\partial \bar{\Psi}_\alpha} = 0 \quad (4)$$

which gives

$$i\hbar\partial_t\Psi_\alpha = -\frac{\hbar^2}{2m_0}(\tau_3 + i\tau_2)\nabla^2\Psi_\alpha + m_0c^2\tau_3\Psi_\alpha \quad (5)$$

where $\alpha = 1, 2$ refers to one of the two components of Ψ . Variation of Ψ gives the conjugate equation

$$-i\hbar\partial_t\bar{\Psi}_\alpha = -\frac{\hbar^2}{2m_0}(\tau_3 + i\tau_2)\nabla^2\bar{\Psi}_\alpha + m_0c^2\tau_3\bar{\Psi}_\alpha \quad (6)$$

These two equations are both forms of the Schrödinger representation of the Klein-Gordon equation, which is

$$\left(i\hbar\frac{\partial}{\partial t} - H_f\right)\Psi = 0 \quad (7)$$

with

$$H_f \equiv (\tau_3 + i\tau_2)\frac{\hbar^2\nabla^2}{2m_0} + \tau_3m_0c^2 \quad (8)$$

We've seen that an integral of the form

$$\langle L \rangle = \int \Psi^\dagger \tau_3 L \Psi d^3x \quad (9)$$

is real if the operator L satisfies the generalized hermiticity condition

$$L^H \equiv \tau_3 L^\dagger \tau_3 = L \quad (10)$$

In eqn 1.111, Greiner shows that H_f satisfies this condition. The operator $i\hbar\partial_t$ has the same effect as H_f so it is generally hermitian as well. As Greiner shows in Ex. 1.7, the general hermiticity of H_f ensures that the action integral is real.

The energy-momentum tensor is given by

$$T_{\mu\nu} = -g_{\mu\nu}\mathcal{L} + \sum_\sigma \frac{\partial\mathcal{L}}{\partial(\partial_\mu\psi_\sigma)} \frac{\partial\psi_\sigma}{\partial x^\nu} \quad (11)$$

where the index σ ranges over the independent fields. In this case, the fields are Ψ and $\bar{\Psi}$, so we have

$$T_{\mu\nu} = -g_{\mu\nu}\mathcal{L} + \frac{\partial\mathcal{L}}{\partial(\partial_\mu\Psi)}\partial_\nu\Psi + \frac{\partial\mathcal{L}}{\partial(\partial_\mu\bar{\Psi})}\partial_\nu\bar{\Psi} \quad (12)$$

The component T_{00} gives the energy density, and Greiner shows in eqns (4) and (5) that the total energy is given by

Note that the 2×2 matrices τ_3 and $i\tau_2$ are both real, so their complex conjugates give the same matrix.

$$E = \int T_{00} d^3x \quad (13)$$

$$= \int \Psi^\dagger \tau_3 H_f \Psi d^3x \quad (14)$$

$$= \langle H_f \rangle \quad (15)$$

where the last line follows from the definition of the mean value of an operator in 9.