

DIELECTRIC CONSTANT

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: Griffiths, David J. (2007) Introduction to Electrodynamics, 3rd Edition; Prentice Hall - Section 4.4.1 and Problem 4.18.

Experimentally, the dipole moment of an atom is proportional to the applied electric field (for small fields). Since the polarization of a dielectric is due to individual atoms within the dielectric being given dipole moments, it should come as no surprise that the polarization density \mathbf{P} of a dielectric is also proportional to the applied field. The relation is written as

$$\mathbf{P} = \epsilon_0 \chi_e \mathbf{E} \quad (1)$$

where χ_e is the *electric susceptibility*. Due to the presence of the ϵ_0 , χ_e is dimensionless.

Not all substances obey such a simple law, but for those that do, they are called *linear dielectrics*.

If this condition holds, we get a simple relationship between the displacement, the field and the polarization. We have

$$\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P} \quad (2)$$

$$= \epsilon_0 (1 + \chi_e) \mathbf{E} \quad (3)$$

The quantity $1 + \chi_e$ is called the *dielectric constant* for a material. The entire proportionality constant is called the *permittivity* ϵ of the material:

$$\epsilon = \epsilon_0 (1 + \chi_e) \quad (4)$$

Since a vacuum cannot be polarized at all, $\chi_e = 0$ for a vacuum, so that $\epsilon = \epsilon_0$ in that case. This is why ϵ_0 is called the *permittivity of free space*.

As a simple example of how these relations can be used, suppose we have a parallel plate capacitor whose plates are separated by a distance $2a$. On one plate there is a surface charge density of σ and on the other there is $-\sigma$. Between the plates are two slabs of dielectric, each of thickness a . Slab 1 (next to the positive plate) has a dielectric constant of 2 and slab 2 has a dielectric constant of 1.5.

We can begin by finding the displacement \mathbf{D} . Using Gauss's law for displacement, we can build a little cylindrical Gaussian pillbox of radius r

with one end in the positive plate and the other in slab 1. By symmetry \mathbf{D} is parallel to the axis of the cylinder so there are no contributions to $\mathbf{D} \cdot d\mathbf{a}$ from the sides of the cylinder. The end of the cylinder inside the plate will have $\mathbf{D} = 0$ (since we're inside a conductor), while the other end will contribute $\pi r^2 D$. The charge enclosed by the cylinder is $\pi r^2 \sigma$ so we get

$$\int \mathbf{D} \cdot d\mathbf{a} = Q \quad (5)$$

$$\pi r^2 D = \pi r^2 \sigma \quad (6)$$

$$D = \sigma \quad (7)$$

This result is independent of the dielectric, so it holds everywhere between the plates. If we assume the positive plate lies above the negative one, we have $\mathbf{D} = -\sigma \hat{\mathbf{z}}$ since the displacement vector points from positive to negative.

From the relation above, we can find \mathbf{E} . For the two slabs, we have

$$\mathbf{E}_1 = \frac{1}{\epsilon_0(1 + \chi_e)_1} \mathbf{D} \quad (8)$$

$$= -\frac{\sigma}{2\epsilon_0} \hat{\mathbf{z}} \quad (9)$$

$$\mathbf{E}_2 = -\frac{\sigma}{1.5\epsilon_0} \hat{\mathbf{z}} \quad (10)$$

The polarization within the two slabs can now be found from 1.

$$\mathbf{P}_1 = \epsilon_0(2 - 1)\mathbf{E}_1 \quad (11)$$

$$= -\frac{\sigma}{2} \hat{\mathbf{z}} \quad (12)$$

$$\mathbf{P}_2 = -\frac{\sigma}{3} \hat{\mathbf{z}} \quad (13)$$

The potential difference between the plates is

$$V = E_1 a + E_2 a \quad (14)$$

$$= \frac{7 a \sigma}{6 \epsilon_0} \quad (15)$$

The bound charges resulting from the polarization can be found. Since the polarization is uniform within each slab, there is no volume bound charge since $\nabla \cdot \mathbf{P} = 0$ everywhere. The surface bound charge is found from

$$\sigma_b = \mathbf{P} \cdot \hat{\mathbf{n}} \quad (16)$$

Applying this at the boundary between the positive plate and slab 1, we get $\sigma_b = -\sigma/2$. At the boundary between slabs 1 and 2, there is a bound charge of $+\sigma/2$ on slab 1 and $-\sigma/3$ on slab 2 (for a net bound charge of $+\sigma/6$ at the boundary). At the boundary between slab 2 and the negative plate, we have $+\sigma/3$.

We can use these bound charges to check the values for the field found above. Choosing a Gaussian cylinder with one end in the positive plate and the other in slab 1, the enclosed charge is $\pi r^2(\sigma - \sigma/2)$ which is equal to ϵ_0 times the surface integral of the field over the cylinder's end cap which is $\pi r^2 E_1$. Thus $E_1 = \sigma/2\epsilon_0$ (pointing downwards) as before. Doing the same calculation for slab 2 we get $E_2 = 2\sigma/3\epsilon_0$ as before.

PINGBACKS

- Pingback: Dielectric examples
- Pingback: Dielectric sphere with free charge
- Pingback: Coaxial cable with dielectric
- Pingback: Dielectric cylinder in uniform electric field
- Pingback: Dielectric sphere in uniform electric field
- Pingback: Point charge embedded in dielectric plane
- Pingback: Point charge in dielectric sphere
- Pingback: Uniqueness of potential in dielectrics
- Pingback: Transmission lines
- Pingback: Skin depth of electromagnetic waves in conductors
- Pingback: [http://physicspages.com/pdf/Griffiths EM/Griffiths Problems 09.23.pdf](http://physicspages.com/pdf/Griffiths%20EM/Griffiths%20Problems%2009.23.pdf)
- Pingback: Frequency dependence of electric permittivity
- Pingback: Microwave shielding for perfect transmission
- Pingback: Maxwell's equations in matter: boundary conditions