

## MAGNETIC VECTOR POTENTIAL OF CONSTANT FIELD

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: Griffiths, David J. (2007) Introduction to Electrodynamics, 3rd Edition; Prentice Hall - Problem 5.24.

The magnetic vector potential  $\mathbf{A}$  for a uniform (that is, constant) magnetic field  $\mathbf{B}$  is

$$\mathbf{A} = -\frac{1}{2}\mathbf{r} \times \mathbf{B} \quad (1)$$

To check this, we calculate its divergence and curl. For the divergence, using a vector calculus identity

$$\nabla \cdot \mathbf{A} = -\frac{1}{2}[\mathbf{B} \cdot (\nabla \times \mathbf{r}) - \mathbf{r} \cdot (\nabla \times \mathbf{B})] \quad (2)$$

The last term is zero since  $\mathbf{B}$  is a constant, and  $\nabla \times \mathbf{r} = 0$  as can be checked by direct calculation, so  $\nabla \cdot \mathbf{A} = 0$  which is one condition required of  $\mathbf{A}$ .

For the curl, we get (omitting terms involving a derivative of  $\mathbf{B}$ , which are all zero):

$$\nabla \times \mathbf{A} = -\frac{1}{2}\nabla \times (\mathbf{r} \times \mathbf{B}) \quad (3)$$

$$-\frac{1}{2}[(\mathbf{B} \cdot \nabla)\mathbf{r} - \mathbf{B}(\nabla \cdot \mathbf{r})] \quad (4)$$

$$= -\frac{1}{2}[\mathbf{B} - 3\mathbf{B}] \quad (5)$$

$$= \mathbf{B} \quad (6)$$

To get the third line from the second, here's a sample calculation of the  $x$  component of the first term:

$$(\mathbf{B} \cdot \nabla)x\hat{\mathbf{x}} = (B_x\partial_x + B_y\partial_y + B_z\partial_z)x\hat{\mathbf{x}} \quad (7)$$

$$= B_x\hat{\mathbf{x}} \quad (8)$$

The  $y$  and  $z$  components work the same way. We've also used  $\nabla \cdot \mathbf{r} = 3$  as can be checked by direct calculation.

Thus the div and curl of  $\mathbf{A}$  give the correct values. It's not unique, since we can add any vector field to  $\mathbf{A}$  so long as its div and curl are both zero. This is true of any constant field, so in general

$$\mathbf{A} = -\frac{1}{2}\mathbf{r} \times \mathbf{B} + \mathbf{C} \quad (9)$$

where  $\mathbf{C}$  is constant over all space.

#### PINGBACKS

Pingback: [Magnetic vector potential from magnetic field](#)

Pingback: [Magnetic vector potential as the curl of another function](#)