

INDUCTANCE OF A COAXIAL CABLE

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Problem 7.28.

Here's another simple example of calculating the energy in the magnetic field and using this to calculate the inductance. We have a coaxial cable with an inner cable of radius R in which the current I flows uniformly, and an outer sheath at radius R in which the current makes its return journey. The magnetic field within the inner cable is given by Ampère's law if we take a loop of radius r centred on the axis. The enclosed current is then

$$I_{enc} = I \frac{r^2}{R^2} = \frac{2\pi r}{\mu_0} B \quad (1)$$

$$B = \frac{\mu_0 r I}{2\pi R^2} \quad (2)$$

Outside the cable, $B = 0$, so the energy per unit length is

$$W_B = \frac{1}{2\mu_0} \int_0^R B^2 2\pi r dr \quad (3)$$

$$= \frac{2\pi}{2\mu_0} \left(\frac{\mu_0 I}{2\pi R^2} \right)^2 \int_0^R r^3 dr \quad (4)$$

$$= \frac{\mu_0 I^2}{16\pi} \quad (5)$$

From $W_B = \frac{1}{2} L I^2$ we get the inductance per unit length:

$$L = \frac{\mu_0}{8\pi} \quad (6)$$

In this case, the inductance is independent of R .