

CHANGING A PARTICLE'S SPEED IN A CYCLOTRON

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Problem 7.48.

We can now revisit the problem of a charged particle in a cyclotron field. Suppose we start with a charged particle of mass m and charge q at rest a distance R from the axis of the cyclotron, and we want to increase the speed of the particle in its circular orbit while keeping it at the same radius (such a device is called a *betatron*). We can do this by varying the magnetic field $\mathbf{B}(t)$ such that the cyclotron relation is always satisfied:

$$(0.1) \quad qvB = \frac{mv^2}{R}$$

$$(0.2) \quad B = \frac{mv}{qR}$$

Taking the time derivative, we get

$$(0.3) \quad \dot{B} = \frac{m\dot{v}}{qR}$$

The changing magnetic field induces a circumferential electric field \mathbf{E} , and since this field is parallel to the particle's direction of motion it will act as the force that accelerates the particle. From Newton's law, $F = m\dot{v} = qE$, so

$$(0.4) \quad \dot{B} = \frac{E}{R}$$

If we assume the cyclotron has cylindrical symmetry, then we can integrate this equation along the particle's orbit, along which both fields are constant in magnitude. That is

$$(0.5) \quad \oint \dot{B}_c dl = \oint \frac{E}{R} dl$$

$$(0.6) \quad \oint E dl = R \oint \dot{B}_c dl$$

$$(0.7) \quad = 2\pi R^2 \dot{B}_c$$

This equation applies on the circumference of the orbit only (not in the interior of the orbit), so we've added a suffix c to B_c to emphasize this point.

From Faraday's law in integral form, we have also that the integral of the electric field is given by the change in flux:

$$(0.8) \quad \oint E d\ell = -\frac{d\Phi}{dt}$$

$$(0.9) \quad = -\int \dot{\mathbf{B}} \cdot d\mathbf{a}$$

where now we are integrating over all points within the orbit.

If we start with the particle at rest in zero field, then we have

$$(0.10) \quad 2\pi R^2 \dot{B}_c = -\int \dot{\mathbf{B}} \cdot d\mathbf{a} = \pi R^2 \dot{\bar{B}}$$

where \bar{B} is the average field across the orbit.

A word about the signs here. Suppose the magnetic field points in the $-z$ direction (as shown in Fig. 7.52 in Griffiths), and we take the area vector to point in the $+z$ direction. Further, if the magnetic field is increasing in magnitude, then $\dot{\mathbf{B}}$ points towards $-z$ as well. Thus $\dot{\mathbf{B}} \cdot d\mathbf{a} < 0$, giving the sign shown.

We can integrate both sides to some time t to get

$$(0.11) \quad \dot{B}_c(t) = \frac{1}{2} \dot{\bar{B}}(t)$$

Thus we can speed up (or slow down, by decreasing the field) the particle by keeping the average field equal to twice the field at the radius of the orbit.