

DUALITY TRANSFORMATION FOR MAGNETIC AND ELECTRIC CHARGE

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Chapter 7, Post 60.

Continuing our excursion in the fantasy world where magnetic monopoles exist, we can explore the *duality transformations*, which describe a rotation in an E-B space, as follows:

$$\mathbf{E}' = \mathbf{E} \cos \alpha + c\mathbf{B} \sin \alpha \quad (1)$$

$$c\mathbf{B}' = c\mathbf{B} \cos \alpha - \mathbf{E} \sin \alpha \quad (2)$$

$$cq'_e = cq_e \cos \alpha + q_m \sin \alpha \quad (3)$$

$$q'_m = q_m \cos \alpha - cq_e \sin \alpha \quad (4)$$

where α is an arbitrary angle and $c = 1/\sqrt{\mu_0\epsilon_0}$. Currents and charge densities transform the same way as the corresponding point charges.

It's a straightforward, though tedious, exercise to verify that Maxwell's equations (including magnetic charge) are invariant under these transformations. The two equations that are different are:

$$\nabla \cdot \mathbf{B} = \mu_0 \rho_m \quad (5)$$

$$\nabla \times \mathbf{E} = -\mu_0 \mathbf{J}_m - \frac{\partial \mathbf{B}}{\partial t} \quad (6)$$

A few sample calculations should show how it goes. We'll do one involving a divergence:

$$c\nabla \cdot \mathbf{B}' = c\nabla \cdot \mathbf{B} \cos \alpha - \nabla \cdot \mathbf{E} \sin \alpha \quad (7)$$

$$= c\mu_0 \rho_m \cos \alpha - \frac{\rho_e}{\epsilon_0} \sin \alpha \quad (8)$$

$$= c\mu_0 (\rho_m \cos \alpha - c\rho_e \sin \alpha) \quad (9)$$

$$= c\mu_0 \rho'_m \quad (10)$$

And another involving a curl:

$$\nabla \times \mathbf{E}' = \nabla \times \mathbf{E} \cos \alpha + c \nabla \times \mathbf{B} \sin \alpha \quad (11)$$

$$= \left(-\mu_0 \mathbf{J}_m - \frac{\partial \mathbf{B}}{\partial t} \right) \cos \alpha + c \left(\mu_0 \mathbf{J}_e + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} \right) \sin \alpha \quad (12)$$

$$= -\mu_0 (\mathbf{J}_m \cos \alpha - c \mathbf{J}_e \sin \alpha) - \frac{\partial \mathbf{B}}{\partial t} \cos \alpha + \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} \sin \alpha \quad (13)$$

$$= -\mu_0 \mathbf{J}'_m - \frac{\partial \mathbf{B}'}{\partial t} \quad (14)$$

The other two equations transform similarly.

If we take the force law for electric and magnetic charges to be

$$\mathbf{F} = q_e (\mathbf{E} + \mathbf{v} \times \mathbf{B}) + q_m \left(\mathbf{B} - \frac{1}{c^2} \mathbf{v} \times \mathbf{E} \right) \quad (15)$$

then the transformed law is

$$\mathbf{F}' = q'_e (\mathbf{E}' + \mathbf{v} \times \mathbf{B}') + q'_m \left(\mathbf{B}' - \frac{1}{c^2} \mathbf{v} \times \mathbf{E}' \right) \quad (16)$$

$$= \left(q_e \cos \alpha + \frac{q_m}{c} \sin \alpha \right) \times \quad (17)$$

$$\left[\mathbf{E} \cos \alpha + c \mathbf{B} \sin \alpha + \mathbf{v} \times \left(\mathbf{B} \cos \alpha - \frac{1}{c} \mathbf{E} \sin \alpha \right) \right] + \quad (18)$$

$$(q_m \cos \alpha - c q_e \sin \alpha) \times \quad (19)$$

$$\left[\mathbf{B} \cos \alpha - \frac{1}{c} \mathbf{E} \sin \alpha - \frac{1}{c^2} \mathbf{v} \times (\mathbf{E} \cos \alpha + c \mathbf{B} \sin \alpha) \right] \quad (20)$$

$$= q_e (\mathbf{E} + \mathbf{v} \times \mathbf{B}) + q_m \left(\mathbf{B} - \frac{1}{c^2} \mathbf{v} \times \mathbf{E} \right) \quad (21)$$

$$= \mathbf{F} \quad (22)$$

To get the penultimate line it is just a matter of multiplying out the first expression and using $\cos^2 \alpha + \sin^2 \alpha = 1$, and cancelling off terms.

PINGBACKS

Pingback: Magnetic monopole force law

Pingback: Radiation from a magnetic dipole composed of monopoles

Pingback: Fields of a moving magnetic monopole