

TRANSMISSION COEFFICIENTS FROM WATER THROUGH GLASS INTO AIR

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Problem 9.36.

Here's another simple example of the transmission coefficient for waves passing from medium 1 through medium 2 into medium 3. Suppose we have an aquarium filled with water (index of refraction $n_1 = \frac{4}{3}$). Light from the aquarium passes (at normal incidence) through a sheet of glass ($n_2 = \frac{3}{2}$) and into air ($n_3 = 1$).

The transmission coefficient is given by

$$(1) \quad T^{-1} = \frac{1}{4n_1n_3} \left[(n_1 + n_3)^2 + \sin^2 \left(\frac{n_2 \omega d}{c} \right) \frac{(n_1^2 - n_2^2)(n_3^2 - n_2^2)}{n_2^2} \right]$$

The minimum value of T occurs when the sine is 1:

$$(2) \quad T_{min}^{-1} = \frac{1}{4n_1n_3} \left[(n_1 + n_3)^2 + \frac{(n_1^2 - n_2^2)(n_3^2 - n_2^2)}{n_2^2} \right]$$

$$(3) \quad T_{min} = 0.935$$

The maximum is when the sine is zero:

$$(4) \quad T_{max} = \frac{4n_1n_3}{(n_1 + n_3)^2}$$

$$(5) \quad = 0.980$$

It doesn't matter much what the frequency of the light is or how thick the glass is; most of the light makes it through in any case. Since 1 is symmetric in n_1 and n_3 a fish inside the aquarium can see out as easily as we can see in.