

GAUGE TRANSFORMATIONS IN ELECTRODYNAMICS

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Problem 10.5.

The potentials V and \mathbf{A} that give rise to a particular configuration of fields \mathbf{E} and \mathbf{B} are not unique. For example, for a point charge at the origin, the usual potentials would be

$$(0.1) \quad V = \frac{q}{4\pi\epsilon_0 r}$$

$$(0.2) \quad \mathbf{A} = 0$$

but as we've seen, the potentials

$$(0.3) \quad V(\mathbf{r}, t) = 0$$

$$(0.4) \quad \mathbf{A}(\mathbf{r}, t) = -\frac{1}{4\pi\epsilon_0} \frac{qt}{r^2} \hat{\mathbf{r}}$$

give rise to the same fields. The question arises: what is the most general set of potentials that give a particular configuration of fields? That is, what changes can we make to V and \mathbf{A} without changing \mathbf{E} and \mathbf{B} ?

The fields can be calculated from the potentials via the equations

$$(0.5) \quad \mathbf{B} = \nabla \times \mathbf{A}$$

$$(0.6) \quad \mathbf{E} = -\nabla V - \frac{\partial \mathbf{A}}{\partial t}$$

We can't multiply V and \mathbf{A} by anything and leave the fields unchanged, but we might be able to add something. Suppose we add a vector function $\boldsymbol{\alpha}(\mathbf{r}, t)$ to \mathbf{A} and a scalar function $\beta(\mathbf{r}, t)$ to V :

$$(0.7) \quad \mathbf{A}' = \mathbf{A} + \boldsymbol{\alpha}$$

$$(0.8) \quad V' = V + \beta$$

To get the same magnetic field, we must have

$$\begin{aligned}
(0.9) \quad \mathbf{B} &= \nabla \times \mathbf{A} \\
(0.10) \quad &= \nabla \times \mathbf{A}' \\
(0.11) \quad &= \nabla \times (\mathbf{A} + \boldsymbol{\alpha}) \\
(0.12) \quad \nabla \times \boldsymbol{\alpha} &= 0
\end{aligned}$$

Since its curl is zero, we can write $\boldsymbol{\alpha}$ as the gradient of a scalar field:

$$(0.13) \quad \boldsymbol{\alpha}(\mathbf{r}, t) = \nabla \lambda'(\mathbf{r}, t)$$

We also have to get the same electric field \mathbf{E} so

$$(0.14) \quad \mathbf{E} = -\nabla V - \frac{\partial \mathbf{A}}{\partial t}$$

$$(0.15) \quad = -\nabla(V + \beta) - \frac{\partial(\mathbf{A} + \boldsymbol{\alpha})}{\partial t}$$

$$(0.16) \quad \nabla \beta + \frac{\partial \boldsymbol{\alpha}}{\partial t} = 0$$

Combining these results, we get

$$(0.17) \quad \nabla \left(\beta + \frac{\partial \lambda'}{\partial t} \right) = 0$$

The term in parentheses must not depend on position (since its gradient is identically zero everywhere), but it might depend on time. Define this term to be $k(t)$; then

$$(0.18) \quad \beta = k(t) - \frac{\partial \lambda'}{\partial t}$$

If we define

$$(0.19) \quad \lambda \equiv \lambda' - \int_0^t k(t') dt'$$

then

$$(0.20) \quad \beta = -\frac{\partial \lambda}{\partial t}$$

Since λ and λ' differ only by a function of time (and not of space), their gradients are equal, so we can use λ in place of λ' in 0.13 without changing

α . Therefore we can change the potentials as follows, without changing the fields they produce:

$$(0.21) \quad \mathbf{A}' = \mathbf{A} + \nabla\lambda$$

$$(0.22) \quad V' = V - \frac{\partial\lambda}{\partial t}$$

where $\lambda = \lambda(\mathbf{r}, t)$ is an arbitrary scalar field. Such a change to the potentials is called a *gauge transformation*.

Example. Earlier, we saw the unusual potentials 0.3 and 0.4 for a point charge at the origin. We can transform them using the gauge function

$$(0.23) \quad \lambda = -\frac{qt}{4\pi\epsilon_0 r}$$

We get

$$(0.24) \quad \nabla\lambda = \frac{qt}{4\pi\epsilon_0 r^2} \hat{\mathbf{r}}$$

$$(0.25) \quad \frac{\partial\lambda}{\partial t} = -\frac{q}{4\pi\epsilon_0 r}$$

so the new potentials are

$$(0.26) \quad V = \frac{q}{4\pi\epsilon_0 r}$$

$$(0.27) \quad \mathbf{A} = 0$$

which is the more usual set of potentials 0.1 and 0.2 for a point charge.

PINGBACKS

Pingback: Coulomb and Lorenz gauges

Pingback: Relativistic electromagnetic potentials