

## GAUGE TRANSFORMATIONS IN ELECTRODYNAMICS

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Problem 10.5.

The potentials  $V$  and  $\mathbf{A}$  that give rise to a particular configuration of fields  $\mathbf{E}$  and  $\mathbf{B}$  are not unique. For example, for a point charge at the origin, the usual potentials would be

$$(1) \quad V = \frac{q}{4\pi\epsilon_0 r}$$

$$(2) \quad \mathbf{A} = 0$$

but as we've seen, the potentials

$$(3) \quad V(\mathbf{r}, t) = 0$$

$$(4) \quad \mathbf{A}(\mathbf{r}, t) = -\frac{1}{4\pi\epsilon_0} \frac{qt}{r^2} \hat{\mathbf{r}}$$

give rise to the same fields. The question arises: what is the most general set of potentials that give a particular configuration of fields? That is, what changes can we make to  $V$  and  $\mathbf{A}$  without changing  $\mathbf{E}$  and  $\mathbf{B}$ ?

The fields can be calculated from the potentials via the equations

$$(5) \quad \mathbf{B} = \nabla \times \mathbf{A}$$

$$(6) \quad \mathbf{E} = -\nabla V - \frac{\partial \mathbf{A}}{\partial t}$$

We can't multiply  $V$  and  $\mathbf{A}$  by anything and leave the fields unchanged, but we might be able to add something. Suppose we add a vector function  $\boldsymbol{\alpha}(\mathbf{r}, t)$  to  $\mathbf{A}$  and a scalar function  $\beta(\mathbf{r}, t)$  to  $V$ :

$$(7) \quad \mathbf{A}' = \mathbf{A} + \boldsymbol{\alpha}$$

$$(8) \quad V' = V + \beta$$

To get the same magnetic field, we must have

$$\begin{aligned}
(9) \quad & \mathbf{B} = \nabla \times \mathbf{A} \\
(10) \quad & = \nabla \times \mathbf{A}' \\
(11) \quad & = \nabla \times (\mathbf{A} + \boldsymbol{\alpha}) \\
(12) \quad & \nabla \times \boldsymbol{\alpha} = 0
\end{aligned}$$

Since its curl is zero, we can write  $\boldsymbol{\alpha}$  as the gradient of a scalar field:

$$(13) \quad \boldsymbol{\alpha}(\mathbf{r}, t) = \nabla \lambda'(\mathbf{r}, t)$$

We also have to get the same electric field  $\mathbf{E}$  so

$$(14) \quad \mathbf{E} = -\nabla V - \frac{\partial \mathbf{A}}{\partial t}$$

$$(15) \quad = -\nabla(V + \beta) - \frac{\partial(\mathbf{A} + \boldsymbol{\alpha})}{\partial t}$$

$$(16) \quad \nabla \beta + \frac{\partial \boldsymbol{\alpha}}{\partial t} = 0$$

Combining these results, we get

$$(17) \quad \nabla \left( \beta + \frac{\partial \lambda'}{\partial t} \right) = 0$$

The term in parentheses must not depend on position (since its gradient is identically zero everywhere), but it might depend on time. Define this term to be  $k(t)$ ; then

$$(18) \quad \beta = k(t) - \frac{\partial \lambda'}{\partial t}$$

If we define

$$(19) \quad \lambda \equiv \lambda' - \int_0^t k(t') dt'$$

then

$$(20) \quad \beta = -\frac{\partial \lambda}{\partial t}$$

Since  $\lambda$  and  $\lambda'$  differ only by a function of time (and not of space), their gradients are equal, so we can use  $\lambda$  in place of  $\lambda'$  in 13 without changing

$\alpha$ . Therefore we can change the potentials as follows, without changing the fields they produce:

$$(21) \quad \mathbf{A}' = \mathbf{A} + \nabla\lambda$$

$$(22) \quad V' = V - \frac{\partial\lambda}{\partial t}$$

where  $\lambda = \lambda(\mathbf{r}, t)$  is an arbitrary scalar field. Such a change to the potentials is called a *gauge transformation*.

**Example.** Earlier, we saw the unusual potentials 3 and 4 for a point charge at the origin. We can transform them using the gauge function

$$(23) \quad \lambda = -\frac{qt}{4\pi\epsilon_0 r}$$

We get

$$(24) \quad \nabla\lambda = \frac{qt}{4\pi\epsilon_0 r^2} \hat{\mathbf{r}}$$

$$(25) \quad \frac{\partial\lambda}{\partial t} = -\frac{q}{4\pi\epsilon_0 r}$$

so the new potentials are

$$(26) \quad V = \frac{q}{4\pi\epsilon_0 r}$$

$$(27) \quad \mathbf{A} = 0$$

which is the more usual set of potentials 1 and 2 for a point charge.

#### PINGBACKS

Pingback: Coulomb and Lorenz gauges

Pingback: Relativistic electromagnetic potentials