

## JEFIMENKO'S EQUATION FOR TIME-DEPENDENT MAGNETIC FIELD

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Problems 10.12.

Using the retarded potentials, we can find a time-dependent expression for the magnetic field  $\mathbf{B}$  to complement the equation we got earlier for the electric field. The potentials are

$$V(\mathbf{r}, t) = \frac{1}{4\pi\epsilon_0} \int \frac{\rho(\mathbf{r}', t_r)}{d} d^3\mathbf{r}' \quad (1)$$

$$\mathbf{A}(\mathbf{r}, t) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{J}(\mathbf{r}', t_r)}{d} d^3\mathbf{r}' \quad (2)$$

where

$$t_r \equiv t - \frac{d}{c} \quad (3)$$

and

$$d \equiv |\mathbf{r} - \mathbf{r}'| \quad (4)$$

$$= \sqrt{(x-x')^2 + (y-y')^2 + (z-z')^2} \quad (5)$$

$$\hat{\mathbf{d}} = \frac{\mathbf{r} - \mathbf{r}'}{d} \quad (6)$$

The magnetic field is

$$\mathbf{B} = \nabla \times \mathbf{A} \quad (7)$$

$$= \frac{\mu_0}{4\pi} \int \left[ \frac{(\nabla \times \mathbf{J}(\mathbf{r}', t_r))}{d} - \mathbf{J}(\mathbf{r}', t_r) \times \nabla \left( \frac{1}{d} \right) \right] d^3\mathbf{r}' \quad (8)$$

where we've used the identity

$$\nabla \times (f\mathbf{A}) = f(\nabla \times \mathbf{A}) - \mathbf{A} \times \nabla f \quad (9)$$

We have

$$\nabla \left( \frac{1}{d} \right) = -\frac{\hat{\mathbf{d}}}{d^2} \quad (10)$$

To get  $\nabla \times \mathbf{J}$ , it's easiest to look at a single component of the curl, say the  $x$  component:

$$(\nabla \times \mathbf{J})_x = \frac{\partial J_z}{\partial y} - \frac{\partial J_y}{\partial z} \quad (11)$$

Using the chain rule

$$\frac{\partial J_z}{\partial y} = \frac{\partial J_z}{\partial t_r} \frac{\partial t_r}{\partial y} \quad (12)$$

$$= -\frac{1}{c} \frac{\partial J_z}{\partial t} \frac{\partial d}{\partial y} \quad (13)$$

$$= -\frac{1}{c} J_z (\nabla d)_y \quad (14)$$

Therefore

$$(\nabla \times \mathbf{J})_x = -\frac{1}{c} \left( J_z (\nabla d)_y - J_y (\nabla d)_z \right) \quad (15)$$

$$= \frac{1}{c} (\mathbf{J} \times \nabla d)_x \quad (16)$$

The other two components work out the same way so we have

$$\nabla \times \mathbf{J} = \frac{1}{c} \mathbf{J} \times \nabla d \quad (17)$$

$$= \frac{1}{c} \mathbf{J} \times \hat{\mathbf{d}} \quad (18)$$

Putting this back into 8 we get

$$\mathbf{B}(\mathbf{r}, t) = \frac{\mu_0}{4\pi} \int \left[ \frac{\dot{\mathbf{J}}(\mathbf{r}', t_r) \times \hat{\mathbf{d}}}{cd} + \mathbf{J}(\mathbf{r}', t_r) \times \frac{\hat{\mathbf{d}}}{d^2} \right] d^3 \mathbf{r}' \quad (19)$$

This is Jefimenko's equation for the magnetic field. For steady currents,  $\dot{\mathbf{J}} = 0$  and the dependence on  $t_r$  disappears, so we're left with the Biot-Savart law:

$$\mathbf{B}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \mathbf{J}(\mathbf{r}') \times \frac{\hat{\mathbf{d}}}{d^2} d^3 \mathbf{r}' \quad (20)$$

In the special case where the current changes slowly enough that we can approximate it by a first-order Taylor series:

$$\mathbf{J}(\mathbf{r}, t_r) = \mathbf{J}(\mathbf{r}, t) + (t_r - t) \dot{\mathbf{J}}(\mathbf{r}, t) \quad (21)$$

$$= \mathbf{J}(\mathbf{r}, t) - \frac{d}{c} \dot{\mathbf{J}}(\mathbf{r}, t) \quad (22)$$

$$\frac{\partial \mathbf{J}}{\partial t_r} = \dot{\mathbf{J}}(\mathbf{r}, t) \quad (23)$$

we get from 19

$$\mathbf{B}(\mathbf{r}, t) = \frac{\mu_0}{4\pi} \int \left[ \frac{\dot{\mathbf{J}}(\mathbf{r}', t)}{cd} + \frac{1}{d^2} \left( \mathbf{J}(\mathbf{r}', t) - \frac{d}{c} \dot{\mathbf{J}}(\mathbf{r}', t) \right) \right] \times \hat{\mathbf{d}} d^3 \mathbf{r}' \quad (24)$$

$$= \frac{\mu_0}{4\pi} \int \frac{\mathbf{J}(\mathbf{r}', t) \times \hat{\mathbf{d}}}{d^2} d^3 \mathbf{r}' \quad (25)$$

That is, for slowly varying currents, to first order, Jefimenko's equation gives the Biot-Savart law where all currents are evaluated at the current time. We're essentially assuming that the travel time for signals from all locations of the current to the observation point is zero. This was the quasistatic approximation.

#### PINGBACKS

Pingback: Relations among charge, current, potentials and fields

Pingback: Electric dipole time-varying potentials; Lorenz gauge condition