

## FORCE OF POINT CHARGE IN A HYPERBOLIC TRAJECTORY ON A FIXED POINT CHARGE

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Problem 10.24.

We'll now return to the point charge  $q_2$  moving on a hyperbolic trajectory along the  $x$  axis. Its position is given by

$$(0.1) \quad x(t) = \sqrt{b^2 + c^2 t^2}$$

Suppose there is a second charge  $q_1$  fixed at  $x = 0$ . What force does  $q_1$  exert on  $q_2$  at time  $t$ ? If we look at the problem from  $q_2$ 's perspective, it needs to know what influence  $q_1$  has on the point  $x$  that it ( $q_2$ , that is) currently occupies. In order for a signal to reach  $x$  at time  $t$ , it had to leave  $q_1$  at time  $t_r = t - \frac{x}{c}$ . But since  $q_1$  is fixed at the origin, the force that  $q_1$  exerts on any charge  $q_2$  at the point  $x$  is always just given by Coulomb's law without any retarded time, that is

$$(0.2) \quad F_2(t) = \frac{q_1 q_2}{4\pi\epsilon_0 x^2}$$

$$(0.3) \quad = \frac{q_1 q_2}{4\pi\epsilon_0} \frac{1}{b^2 + c^2 t^2}$$

The total impulse delivered to  $q_2$  is

$$(0.4) \quad I_2 = \int_{-\infty}^{\infty} F_2(t) dt$$

$$(0.5) \quad = \frac{q_1 q_2}{4\pi\epsilon_0 b^2} \int_{-\infty}^{\infty} \frac{1}{1 + \frac{c^2}{b^2} t^2} dt$$

$$(0.6) \quad = \frac{q_1 q_2}{4\pi\epsilon_0 bc}$$

To calculate the force  $q_2$  exerts on  $q_1$  we *do* need the retarded time, since  $q_2$  is moving. [Incidentally, it might seem that by considering  $q_1$  as at rest and  $q_2$  as moving, and treating the two cases differently, we're violating the principle of relativity, but we're not. The reason is that  $q_2$  is not in an inertial frame; the hyperbolic trajectory means that its velocity is never constant, so  $q_1$  and  $q_2$  are not equivalent.]

FORCE OF POINT CHARGE IN A HYPERBOLIC TRAJECTORY ON A FIXED POINT CHARGE

The retarded time is calculated from

$$(0.7) \quad |\mathbf{r} - \mathbf{w}(t_r)| = c(t - t_r)$$

Here  $\mathbf{r} = 0$  is the location of  $q_1$  and  $\mathbf{w}(t_r) = \sqrt{b^2 + c^2 t_r^2} \hat{\mathbf{x}}$  is the position of  $q_2$ . We get

$$(0.8) \quad \sqrt{b^2 + c^2 t_r^2} = c(t - t_r)$$

$$(0.9) \quad t_r = \frac{t}{2} - \frac{b^2}{2c^2 t}$$

Note that as  $t \rightarrow 0$ ,  $t_r \rightarrow -\infty$  so  $q_2$  is not visible to  $q_1$  before  $t = 0$ , so the retarded potential is zero for  $t < 0$ . The force for  $t > 0$  is therefore (the minus sign indicates the force is to the left if both charges are the same sign)

$$(0.10) \quad F_1(t) = -\frac{q_1 q_2}{4\pi\epsilon_0} \frac{1}{b^2 + \frac{c^2}{4} \left(t - \frac{b^2}{c^2 t}\right)^2}$$

$$(0.11) \quad = -\frac{q_1 q_2}{\pi\epsilon_0} \frac{c^2 t^2}{4c^2 b^2 t^2 + (c^2 t^2 - b^2)^2}$$

$$(0.12) \quad = -\frac{q_1 q_2}{\pi\epsilon_0} \frac{c^2 t^2}{(c^2 t^2 + b^2)^2}$$

The total impulse is found by integrating  $F_1$  from  $t = 0$  to infinity, since there is no force for  $t < 0$ .

$$(0.13) \quad I_1 = -\frac{q_1 q_2}{\pi\epsilon_0} \int_0^\infty \frac{c^2 t^2}{(c^2 t^2 + b^2)^2} dt$$

$$(0.14) \quad = -\frac{q_1 q_2}{4\pi\epsilon_0 b c}$$

[The integral can be done by parts, although I used Maple.] Thus the two impulses are equal and opposite.