

## RADIATION FROM A CHARGE IN HYPERBOLIC MOTION

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Chapter 11, Post 31.

A particle moving along the  $x$  axis in hyperbolic motion has a position given by

$$(1) \quad x(t) = \sqrt{b^2 + c^2 t^2}$$

where  $b$  is a constant giving the closest approach to the origin. If the particle has charge  $q$  then its radiation is given by

$$(2) \quad P = \frac{\mu_0 q^2 \gamma^6 a^2}{6\pi c}$$

To calculate this we need

$$(3) \quad v(t) = \frac{tc^2}{\sqrt{b^2 + c^2 t^2}} = \frac{tc^2}{x}$$

$$(4) \quad a(t) = \frac{c^2}{x} - \frac{c^4 t^2}{x^3} = \frac{c^2 b^2}{x^3}$$

$$(5) \quad \gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$$

$$(6) \quad = \frac{1}{\sqrt{1 - \frac{t^2 c^2}{x^2}}}$$

$$(7) \quad = \frac{x}{b}$$

The power is therefore

$$(8) \quad P = \frac{\mu_0 q^2}{6\pi c} \left(\frac{x}{b}\right)^6 \left(\frac{c^2 b^2}{x^3}\right)^2$$

$$(9) \quad = \frac{\mu_0 q^2 c^3}{6\pi b^2}$$

Thus the radiated power is constant as the charge moves along its trajectory.

The radiation reaction force is given by

$$(10) \quad F_{rad} = \frac{\mu_0 q^2 \gamma^4}{6\pi c} \left( \dot{a} + \frac{3a^2 \gamma^2 v}{c^2} \right)$$

for which we need

$$(11) \quad \dot{a} = -\frac{3c^4 b^2 t}{x^5}$$

We get

$$(12) \quad \frac{3a^2 \gamma^2 v}{c^2} = \frac{3}{c^2} \left( \frac{c^2 b^2}{x^3} \right)^2 \left( \frac{x}{b} \right)^2 \left( \frac{tc^2}{x} \right)$$

$$(13) \quad = \frac{3c^4 b^2 t}{x^5}$$

$$(14) \quad = -\dot{a}$$

$$(15) \quad F_{rad} = 0$$

This is a rather curious case in that although the charge radiates, there is no radiation reaction force.