

VELOCITY ADDITION FORMULAS FOR ALL 3 DIRECTIONS

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Chapter 12, Problem 12.14.

Reference: Carroll, Bradley W. & Ostlie, Dale A. (2007), *An Introduction to Modern Astrophysics*, 2nd Edition; Pearson Education - Chapter 4, Problem 4.10.

The relativistic velocity addition formula is most easily derived from the Lorentz transformations:

$$\bar{x} = \gamma(x - vt) \quad (1)$$

$$\bar{t} = \gamma\left(t - \frac{xv}{c^2}\right) \quad (2)$$

As usual, system \bar{S} is moving to the right relative to system S with velocity v . If an object is moving in the S system with velocity $u = dx/dt$, then we can take differentials on both sides of the above equations to get

$$d\bar{x} = \gamma(dx - v dt) \quad (3)$$

$$d\bar{t} = \gamma\left(dt - \frac{v}{c^2}dx\right) \quad (4)$$

$$\bar{u} = \frac{d\bar{x}}{d\bar{t}} \quad (5)$$

$$= \frac{dx - v dt}{dt - \frac{v}{c^2}dx} \quad (6)$$

$$= \frac{(dx/dt) - v}{1 - \frac{v}{c^2}(dx/dt)} \quad (7)$$

$$= \frac{u - v}{1 - uv/c^2} \quad (8)$$

$$\dot{\bar{x}} = \frac{\dot{x} - v}{1 - \dot{x}v/c^2} \quad (9)$$

By symmetry (or algebra if you don't trust the physical argument) this relation can be inverted to give

$$u = \frac{\bar{u} + v}{1 + \bar{u}v/c^2} \quad (10)$$

We can also derive the formulas for transforming velocity components perpendicular to the motion. Although perpendicular distances don't change under the Lorentz transformation, perpendicular *velocities* do, because they involve the ratio of perpendicular distance to time, and time *does* change from one system to another. We get, with a dot indicating a time derivative in the respective system:

$$\dot{\bar{y}} = \frac{d\bar{y}}{d\bar{t}} \quad (11)$$

$$= \frac{dy}{\gamma \left(dt - \frac{v}{c^2} dx \right)} \quad (12)$$

$$= \frac{\dot{y}}{\gamma (1 - \dot{x}v/c^2)} \quad (13)$$

$$\dot{\bar{z}} = \frac{\dot{z}}{\gamma (1 - \dot{x}v/c^2)} \quad (14)$$

Thus perpendicular velocities depend not only on the perpendicular velocity in the original system, but also on the horizontal velocity.

Example. Suppose we have a spotlight mounted on the roof of our usual train, and the spotlight is aimed to point towards the rear of the train, making an angle of θ with the roof of the train. If the train moves with speed v , what angle will a ground observer see the beam make with the train's roof?

In the train's system, the light's velocity components are

$$\dot{x} = -c \cos \theta \quad (15)$$

$$\dot{z} = c \sin \theta \quad (16)$$

In the ground system, we get from 10:

$$\dot{\bar{x}} = \frac{-c \cos \theta + v}{1 - v(\cos \theta)/c} \quad (17)$$

$$\dot{\bar{z}} = \frac{c \sin \theta}{\gamma (1 - v(\cos \theta)/c)} \quad (18)$$

$$\tan \bar{\theta} = -\frac{\dot{\bar{z}}}{\dot{\bar{x}}} \quad (19)$$

$$= \frac{c \sin \theta}{\gamma (c \cos \theta - v)} \quad (20)$$

As a check, we can work out $\dot{x}^2 + \dot{z}^2 = c^2$ (after simplifying). When the beam points directly up in the train's frame ($\theta = \frac{\pi}{2}$), it points slightly forward in the ground's frame. When $\cos\theta = \frac{v}{c}$, $\tan\bar{\theta} = \infty$ so the beam points directly up in the ground's frame.

PINGBACKS

Pingback: Velocity addition: chasing space pirates viewed in four reference frames

Pingback: Spacetime diagrams: an example

Pingback: Four-velocity versus ordinary velocity: an example

Pingback: Relativistic momentum and energy

Pingback: Relativistic transformation of force, electric and magnetic fields

Pingback: Transforming the electric field in relativity