

COMPOUND LORENTZ TRANSFORMATIONS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Chapter 12, Post 18.

The Lorentz transformations can be written in matrix form as

$$(0.1) \quad \Lambda_x = \begin{bmatrix} \gamma & -\beta\gamma & 0 & 0 \\ -\beta\gamma & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

where the 0 (ct) component is the first row and first column, followed by the 1, 2, and 3 directions in order. This matrix is for relative motion along the 1 axis.

The Galilean transformations can be written as a matrix as well, where the first coordinate is just t rather than ct :

$$(0.2) \quad \Gamma = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -v & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

or if we want to use the same symbols as in the Lorentz case, where the top row of Γ is a ct coordinate, we can write

$$(0.3) \quad \Gamma = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -\beta & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

The Lorentz transformation along the 2 (y) axis is obtained by putting the transformation terms in row and column 2:

$$(0.4) \quad \Lambda_y = \begin{bmatrix} \gamma & 0 & -\beta\gamma & 0 \\ 0 & 1 & 0 & 0 \\ -\beta\gamma & 0 & \gamma & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

1

If we apply a Lorentz transformation first in the x and then in the y direction (with different relative velocities), we get the compound matrix:

$$(0.5) \quad \Lambda_y \Lambda_x = \begin{bmatrix} \gamma_y & 0 & -\beta_y \gamma_y & 0 \\ 0 & 1 & 0 & 0 \\ -\beta_y \gamma_y & 0 & \gamma_y & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \gamma_x & -\beta_x \gamma_x & 0 & 0 \\ -\beta_x \gamma_x & \gamma_x & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(0.6) \quad = \begin{bmatrix} \gamma_y \gamma_x & -\gamma_y \gamma_x \beta_x & -\beta_y \gamma_y & 0 \\ -\beta_x \gamma_x & \gamma_x & 0 & 0 \\ -\beta_y \gamma_y \gamma_x & \gamma_y \gamma_x \beta_x \beta_y & \gamma_y & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Note that although Λ_x and Λ_y are both symmetric, their product is not. This means that applying the transformations in the opposite order gives a different result.

$$(0.7) \quad \Lambda_x \Lambda_y = \Lambda_x^T \Lambda_y^T$$

$$(0.8) \quad = (\Lambda_y \Lambda_x)^T$$

$$(0.9) \quad = \begin{bmatrix} \gamma_y \gamma_x & -\beta_x \gamma_x & -\beta_y \gamma_y \gamma_x & 0 \\ -\gamma_y \gamma_x \beta_x & \gamma_x & \gamma_y \gamma_x \beta_x \beta_y & 0 \\ -\beta_y \gamma_y & 0 & \gamma_y & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$