

PAIR ANNIHILATION WITH AN ELECTRON AND A POSITRON

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Chapter 12, Post 35.

Here's a slightly more involved example of conservation of energy and momentum in relativity. Suppose an electron with momentum p_e hits a positron (anti-electron) and they annihilate each other (a process known as *pair annihilation*), producing two photons.

In passing, it's worth noting that pair annihilation must produce at least 2 photons. If we looked at the problem in the centre of momentum frame, the total momentum of the system must be zero both before and after the collision. If only one photon were produced, it is impossible for it to have zero momentum.

Before the collision, we have

$$(0.1) \quad E_{tot} = mc^2 + \sqrt{mc^2 + p_e^2 c^2}$$

After the collision, suppose that one photon emerges at $60^\circ = \frac{\pi}{3}$ radians to the incident electron's direction. From conservation of momentum, we have

$$(0.2) \quad p_1 \sin \frac{\pi}{3} = p_2 \sin \theta$$

$$(0.3) \quad p_1 \cos \frac{\pi}{3} + p_2 \cos \theta = p_e$$

where θ is the angle at which photon 2 emerges. We can eliminate θ as follows (using $\sin \frac{\pi}{3} = \frac{\sqrt{3}}{2}$, $\cos \frac{\pi}{3} = \frac{1}{2}$):

$$(0.4) \quad p_2^2 (\sin^2 \theta + \cos^2 \theta) = \frac{3}{4} p_1^2 + \left(p_e - \frac{1}{2} p_1 \right)^2$$

$$(0.5) \quad p_2^2 = p_1^2 - p_1 p_e + p_e^2$$

For a photon, $E = cp$, so the total energy after the collision is

$$(0.6) \quad E_{tot} = cp_1 + cp_2$$

From conservation of energy and 0.1, we get

$$(0.7)$$

$$c^2 p_2^2 = (E_{tot} - cp_1)^2$$

$$(0.8)$$

$$= \left(mc^2 + \sqrt{m^2 c^4 + p_e^2 c^2} - cp_1 \right)^2$$

$$(0.9)$$

$$= \left(mc^2 + \sqrt{m^2 c^4 + p_e^2 c^2} \right)^2 - 2 \left(mc^2 + \sqrt{m^2 c^4 + p_e^2 c^2} \right) cp_1 + c^2 p_1^2$$

We can now use 0.5:

$$(0.10)$$

$$c^2 p_2^2 = c^2 p_1^2 - c^2 p_1 p_e + c^2 p_e^2$$

$$(0.11)$$

$$= \left(mc^2 + \sqrt{m^2 c^4 + p_e^2 c^2} \right)^2 - 2 \left(mc^2 + \sqrt{m^2 c^4 + p_e^2 c^2} \right) cp_1 + c^2 p_1^2$$

Solving for cp_1 we get

$$(0.12) \quad cp_1 = \frac{\left(mc^2 + \sqrt{m^2 c^4 + p_e^2 c^2} \right)^2 - c^2 p_e^2}{2 \left(mc^2 + \sqrt{m^2 c^4 + p_e^2 c^2} \right) - cp_e}$$

$$(0.13) \quad = \frac{2m^2 c^4 + 2mc^2 \sqrt{m^2 c^4 + p_e^2 c^2}}{2 \left(mc^2 + \sqrt{m^2 c^4 + p_e^2 c^2} \right) - cp_e}$$

$$(0.14) \quad = \frac{mc^2}{1 - cp_e / \left[2 \left(mc^2 + \sqrt{m^2 c^4 + p_e^2 c^2} \right) \right]}$$

Thus photon 1 has an energy greater than the rest mass of the electron. Note that if the pair annihilation occurs with both the electron and positron at rest ($p_e = 0$), then $cp_1 = cp_2 = mc^2$ as we'd expect.