

MOTION UNDER A CONSTANT MINKOWSKI FORCE

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

Reference: Griffiths, David J. (2007), Introduction to Electrodynamics, 3rd Edition; Pearson Education - Chapter 12, Problem 12.60.

The Minkowski force \mathbf{K} is the rate of change of four-momentum with respect to proper time, and allows Newton's law to be written in its natural form

$$\mathbf{K} = m\alpha \quad (1)$$

where α is the proper acceleration, or second derivative of position with respect to proper time. Here we'll investigate the behaviour of a particle subject to a constant Minkowski force in one dimension.

In terms of ordinary force, we have

$$K = \frac{dp}{d\tau} = \frac{dp}{dt} \frac{dt}{d\tau} = \frac{1}{\sqrt{1-u^2/c^2}} F \quad (2)$$

The ordinary momentum p is

$$p = \frac{mu}{\sqrt{1-u^2/c^2}} \quad (3)$$

so its derivative is

$$\frac{dp}{dt} = \frac{m}{\sqrt{1-u^2/c^2}} \frac{du}{dt} + \frac{mu^2}{(1-u^2/c^2)^{3/2}} \frac{du}{dt} \quad (4)$$

Inserting this into 2 we get

$$\frac{K}{m} dt = \frac{du}{1-u^2/c^2} + \frac{u^2 du}{(1-u^2/c^2)^2} \quad (5)$$

We can integrate both sides (using software, or integral tables) to get

$$\frac{K}{m} t + C = \frac{c}{4} \ln \left[\frac{c+u}{c-u} \right] + \frac{c^2}{4} \left[\frac{1}{c-u} - \frac{1}{c+u} \right] \quad (6)$$

where C is a constant of integration. If the initial conditions are $u = 0$ at $t = 0$, then $C = 0$ and we have

$$\frac{K}{m}t = \frac{c}{4} \ln \left[\frac{c+u}{c-u} \right] + \frac{c^2}{4} \left[\frac{1}{c-u} - \frac{1}{c+u} \right] \quad (7)$$

This is an implicit equation for the speed of the particle as a function of time. If we want the position as a function of time, we need a relation between u and x . Returning to 2 and 3 we have

$$\sqrt{1-u^2/c^2} \frac{K}{m} = \frac{d}{dt} \left(\frac{u}{\sqrt{1-u^2/c^2}} \right) \quad (8)$$

We can use the chain rule to convert the derivative on the RHS to a derivative with respect to x by multiplying both sides by dt/dx

$$\frac{dt}{dx} \sqrt{1-u^2/c^2} \frac{K}{m} = \frac{dt}{dx} \frac{d}{dt} \left(\frac{u}{\sqrt{1-u^2/c^2}} \right) = \frac{d}{dx} \left(\frac{u}{\sqrt{1-u^2/c^2}} \right) \quad (9)$$

Now $dx/dt = u$ so $dt/dx = 1/u$ and

$$\frac{\sqrt{1-u^2/c^2}}{u} \frac{K}{m} = \frac{d}{dx} \left(\frac{u}{\sqrt{1-u^2/c^2}} \right) \quad (10)$$

If we call the expression in the parentheses on the RHS A , then we can integrate with respect to x (since K/m is a constant):

$$A \equiv \frac{u}{\sqrt{1-u^2/c^2}} \quad (11)$$

$$\frac{1}{A} \frac{K}{m} = \frac{dA}{dx} \quad (12)$$

$$\frac{K}{m}x + C = \frac{1}{2}A^2 \quad (13)$$

Again, starting from rest at the origin we have $u = 0$ when $x = 0$ so $A = 0$ also, and therefore $C = 0$, so we have

$$A = \frac{u}{\sqrt{1-u^2/c^2}} = \sqrt{\frac{2Kx}{m}} \quad (14)$$

At this point we could get a relation between x and t by solving 14 for u in terms of x and then substituting this into 7. For reference, we get

$$u = \sqrt{\frac{2Kx}{m}} \frac{1}{\sqrt{1+2Kx/mc^2}} \quad (15)$$

so substituting will give something of a mess. To get the answer given in

Griffiths requires a bit of algebra, but here is how I did it. Griffiths defines the quantity z as

$$z \equiv \sqrt{\frac{2Kx}{mc^2}} \quad (16)$$

$$= \frac{A}{c} \quad (17)$$

$$= \frac{u}{c\sqrt{1-u^2/c^2}} \quad (18)$$

The quantities appearing in Griffiths's answer are

$$\sqrt{1+z^2} = \frac{c}{\sqrt{c^2-u^2}} \quad (19)$$

$$z\sqrt{1+z^2} = \frac{u}{c(1-u^2/c^2)} \quad (20)$$

We can rewrite 7 to get

$$\frac{2Kt}{mc} = \frac{1}{2} \ln \left[\frac{c+u}{c-u} \right] + \frac{c}{2} \left[\frac{1}{c-u} - \frac{1}{c+u} \right] \quad (21)$$

We'll deal with the logarithm first. Its argument is

$$\frac{c+u}{c-u} = \frac{(c+u)^2}{c^2(1-u^2/c^2)} \quad (22)$$

$$= \frac{2u}{c(1-u^2/c^2)} + \frac{u^2+c^2}{c^2-u^2} \quad (23)$$

$$= \frac{2u}{c(1-u^2/c^2)} + \frac{c^2-u^2+2u^2}{c^2-u^2} \quad (24)$$

$$= \frac{2u}{c(1-u^2/c^2)} + 1 + \frac{2u^2}{c^2(1-u^2/c^2)} \quad (25)$$

Now we also have

$$\left(z + \sqrt{1+z^2} \right)^2 = 2z^2 + 2z\sqrt{1+z^2} + 1 \quad (26)$$

$$= \frac{2u^2}{c^2(1-u^2/c^2)} + \frac{2u}{c(1-u^2/c^2)} + 1 \quad (27)$$

$$= \frac{c+u}{c-u} \quad (28)$$

Therefore

$$\frac{1}{2} \ln \left[\frac{c+u}{c-u} \right] = \ln \sqrt{\frac{c+u}{c-u}} \quad (29)$$

$$= \ln \left(z + \sqrt{1+z^2} \right) \quad (30)$$

For the second term in 21, we have

$$\frac{c}{2} \left[\frac{1}{c-u} - \frac{1}{c+u} \right] = \frac{c}{2} \frac{2u}{c^2(1-u^2/c^2)} \quad (31)$$

$$= \frac{u}{c(1-u^2/c^2)} \quad (32)$$

$$= z\sqrt{1+z^2} \quad (33)$$

Putting it all together, we have

$$\frac{2Kt}{mc} = \ln \left(z + \sqrt{1+z^2} \right) + z\sqrt{1+z^2} \quad (34)$$