

## BUFFON'S NEEDLE: ESTIMATING PI

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

Post date: 24 Jan 2011.

Reference: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 1.13.

This interesting little problem serves well to illustrate the notion of a probability density and its application to an experiment which can be done at home. It is known as Buffon's needle, since it is believed that Georges-Louis Leclerc, Comte de Buffon, first posed the problem in the 18th century.

Suppose we have a needle of length  $l$  and we drop this needle onto a sheet of paper on which there are a number of parallel lines spaced a distance  $l$  apart. What is the probability that the dropped needle will cross a line?

One way of analyzing this problem is to begin by considering the needle as the hypotenuse of a right triangle, with sides parallel and perpendicular to the parallel lines. Then if the needle makes an angle  $\theta$  with the lines, the sides of this triangle are  $l \cos \theta$  for the parallel side and  $l \sin \theta$  for the perpendicular side. The side of the triangle parallel to the lines is of no interest here; what we are interested in the perpendicular component.

To see this, suppose we placed a needle of length  $x < l$  on the paper in such a way that it was perpendicular to the lines. Such a needle would cover a fraction  $x/l$  of the distance between two adjacent lines. Thus the probability that a line would cross this specially dropped needle is just  $x/l$ . (Note that if  $x = l$  the probability is 1, since such a needle covers the entire distance between the lines.)

Therefore, the probability that a needle that makes an angle  $\theta$  with the lines crosses a line is  $(l \sin \theta)/l = \sin \theta$ .

How likely is the needle to drop at an angle  $\theta$ ? Since  $\theta$  is a continuous variable, we need a probability density, rather than just a simple probability. Assuming that the needle is equally likely to drop at any angle between 0 and  $\pi$  ( $180^\circ$ ), the density must be a constant, and must integrate to 1 over the range of valid  $\theta$ , so it must be  $\rho(\theta) = \frac{1}{\pi}$ .

The probability  $P(\text{crosses line})$  that a randomly dropped needle crosses a line is therefore

$$P(\text{crosses line}) = \int_0^\pi P(\text{angle } \theta \text{ crosses line})\rho(\theta)d\theta \quad (1)$$

$$= \frac{1}{\pi} \int_0^\pi \sin \theta \, d\theta \quad (2)$$

$$= \frac{2}{\pi} \quad (3)$$

$$\approx 0.6366 \quad (4)$$

Besides being another of those curious situations where  $\pi$  pops up unexpectedly, this result offers the possibility of an interesting way of whiling away a rainy afternoon. Simply by dropping a needle repeatedly onto lined paper, you can do an experiment that will determine the value of  $\pi$ , since

$$\pi = \frac{2}{P(\text{crosses line})} \quad (5)$$

$$= \frac{2}{(\text{Fraction of needles crossing lines})} \quad (6)$$

$$= \frac{2(\text{Number of needles dropped})}{\text{Number of needles crossing lines}} \quad (7)$$

OK, you would need to drop a lot of needles to get a decent value of  $\pi$ , but it's interesting that there is such a simple method for getting even an approximate value of  $\pi$  without using circles, triangles, angles, measurement or anything more than just counting.

#### PINGBACKS

Pingback: Continuous probability distribution: needle on a pivot