

BUFFON'S NEEDLE: ESTIMATING PI

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

Reference: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 1.13.

This interesting little problem serves well to illustrate the notion of a probability density and its application to an experiment which can be done at home. It is known as Buffon's needle, since it is believed that Georges-Louis Leclerc, Comte de Buffon, first posed the problem in the 18th century.

Suppose we have a needle of length l and we drop this needle onto a sheet of paper on which there are a number of parallel lines spaced a distance l apart. What is the probability that the dropped needle will cross a line?

One way of analyzing this problem is to begin by considering the needle as the hypotenuse of a right triangle, with sides parallel and perpendicular to the parallel lines. Then if the needle makes an angle θ with the lines, the sides of this triangle are $l \cos \theta$ for the parallel side and $l \sin \theta$ for the perpendicular side. The side of the triangle parallel to the lines is of no interest here; what we are interested in the perpendicular component.

To see this, suppose we placed a needle of length $x < l$ on the paper in such a way that it was perpendicular to the lines. Such a needle would cover a fraction x/l of the distance between two adjacent lines. Thus the probability that a line would cross this specially dropped needle is just x/l . (Note that if $x = l$ the probability is 1, since such a needle covers the entire distance between the lines.)

Therefore, the probability that a needle that makes an angle θ with the lines crosses a line is $(l \sin \theta)/l = \sin \theta$.

How likely is the needle to drop at an angle θ ? Since θ is a continuous variable, we need a probability density, rather than just a simple probability. Assuming that the needle is equally likely to drop at any angle between 0 and π (180°), the density must be a constant, and must integrate to 1 over the range of valid θ , so it must be $\rho(\theta) = \frac{1}{\pi}$.

The probability $P(\text{crosses line})$ that a randomly dropped needle crosses a line is therefore

$$(0.1) \quad P(\text{crosses line}) = \int_0^\pi P(\text{angle } \theta \text{ crosses line}) \rho(\theta) d\theta$$

$$(0.2) \quad = \frac{1}{\pi} \int_0^\pi \sin \theta d\theta$$

$$(0.3) \quad = \frac{2}{\pi}$$

$$(0.4) \quad \approx 0.6366$$

Besides being another of those curious situations where π pops up unexpectedly, this result offers the possibility of an interesting way of whiling away a rainy afternoon. Simply by dropping a needle repeatedly onto lined paper, you can do an experiment that will determine the value of π , since

$$(0.5) \quad \pi = \frac{2}{P(\text{crosses line})}$$

$$(0.6) \quad = \frac{2}{(\text{Fraction of needles crossing lines})}$$

$$(0.7) \quad = \frac{2(\text{Number of needles dropped})}{\text{Number of needles crossing lines}}$$

OK, you would need to drop a lot of needles to get a decent value of π , but it's interesting that there is such a simple method for getting even an approximate value of π without using circles, triangles, angles, measurement or anything more than just counting.

PINGBACKS

Pingback: Continuous probability distribution: needle on a pivot