

HERMITE POLYNOMIALS - GENERATION

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Post date: 21 Jul 2012.

Reference: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 2.16.

In the solution of the Schrödinger equation for the harmonic oscillator, we found that the wave function can be expressed as a power series:

$$\psi(y) = e^{-y^2/2} \sum_{j=0}^{\infty} a_j y^j \quad (1)$$

where y was introduced as a shorthand variable:

$$y \equiv \sqrt{\frac{m\omega}{\hbar}} x \quad (2)$$

and the coefficients a_j satisfy the recursion relation

$$a_{j+2} = \frac{2j+1-\varepsilon}{(j+1)(j+2)} a_j \quad (3)$$

where ε is another shorthand variable for the energy:

$$\varepsilon \equiv \frac{2E}{\hbar\omega} \quad (4)$$

By requiring the recursion relation to terminate at various values of j we can generate the polynomials for the various energy states, which turn out to be the Hermite polynomials.

For example, if we take the highest value of j to be 5, then we must have

$$a_7 = \frac{11-\varepsilon}{42} a_5 \quad (5)$$

$$= 0 \quad (6)$$

$$\varepsilon = 11 \quad (7)$$

$$E = \frac{11}{2} \hbar\omega \quad (8)$$

To get the coefficients, we can start with $a_0 = 0$ (since all even terms must be zero if we want $a_5 \neq 0$) and $a_1 = 1$. Then we get

$$a_{j+2} = \frac{2j-10}{(j+1)(j+2)}a_j \quad (9)$$

so $a_3 = -\frac{4}{3}$, $a_5 = \left(-\frac{1}{5}\right)\left(-\frac{4}{3}\right) = \frac{4}{15}$. The Hermite polynomial is

$$H_5(x) = A_5 \left[\frac{4}{15}x^5 - \frac{4}{3}x^3 + x \right] \quad (10)$$

where A_5 is a constant that is set by convention to make the coefficients satisfy some specified rule.

If we require the coefficient of the highest power n to be 2^n we can multiply this polynomial by 120 to get

$$H_5(x) = 32x^5 - 160x^3 + 120x \quad (11)$$

For H_6 , all the odd terms are zero, and we require $a_8 = 0$, so we get

$$\varepsilon = 13 \quad (12)$$

$$E = \frac{13}{2}\hbar\omega \quad (13)$$

$$a_{j+2} = \frac{2j-12}{(j+1)(j+2)}a_j \quad (14)$$

Taking $a_0 = 1$ and $a_1 = 0$, we get $a_2 = -6$, $a_4 = 4$, $a_6 = -\frac{8}{15}$. Requiring the coefficient of a_6 to be $2^6 = 64$ we get

$$H_6(x) = 64x^6 - 480x^4 + 720x^2 - 120 \quad (15)$$

PINGBACKS

Pingback: Harmonic oscillator - series solution

Pingback: Harmonic oscillator: Hermite polynomials and orthogonality of eigenfunctions