

COMPLEX EXPONENTIALS AND TRIG FUNCTIONS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

Reference: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 2.18.

Solutions to the Schrödinger equation are written either as complex exponentials or a sum of sine and cosine functions. For example, we used the former in the solution for the free particle and the latter in the infinite square well. In fact these two ways are equivalent for writing a function, as we can demonstrate.

We start with

$$(0.1) \quad Ae^{ikx} + Be^{-ikx}$$

and we want to write this as $C \cos kx + D \sin kx$.

We can expand the exponentials in terms of sine and cosine to get

$$(0.2) \quad Ae^{ikx} + Be^{-ikx} = (A+B) \cos kx + i(A-B) \sin kx$$

$$(0.3) \quad \equiv C \cos kx + D \sin kx$$

So

$$(0.4) \quad C = A+B$$

$$(0.5) \quad D = i(A-B)$$

or, the other way round:

$$(0.6) \quad A = \frac{1}{2}(C - iD)$$

$$(0.7) \quad B = \frac{1}{2}(C + iD)$$

PINGBACKS

Pingback: [The free particle](#)