

## HERMITIAN OPERATORS - EQUIVALENCE OF CONDITIONS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 3.3.

We've seen that a Hermitian operator  $Q$  satisfies the condition

$$(0.1) \quad \int_a^b g^* Qf \, dx = \int_a^b f(Qg)^* \, dx$$

where  $f$  and  $g$  are any functions in Hilbert space that satisfy certain boundary conditions at  $a$  and  $b$  (see earlier post for a full discussion).

Using the more compact bracket notation, this condition can be written as

$$(0.2) \quad \langle g | \hat{Q}f \rangle = \langle \hat{Q}g | f \rangle$$

It is also possible to arrive at this result if we start off with the apparently less restrictive condition

$$(0.3) \quad \langle h | \hat{Q}h \rangle = \langle \hat{Q}h | h \rangle$$

where  $h$  is any function in Hilbert space.

To show this, we start with  $h = f + g$ :

$$(0.4) \quad \langle h | \hat{Q}h \rangle = \langle f + g | \hat{Q}f + \hat{Q}g \rangle$$

$$(0.5) \quad = \langle f | \hat{Q}f \rangle + \langle g | \hat{Q}g \rangle + \langle f | \hat{Q}g \rangle + \langle g | \hat{Q}f \rangle$$

But we are assuming that  $\langle h | \hat{Q}h \rangle = \langle \hat{Q}h | h \rangle$  so we can write this equation as:

$$(0.6) \quad \langle \hat{Q}h | h \rangle = \langle \hat{Q}f + \hat{Q}g | f + g \rangle$$

$$(0.7) \quad = \langle \hat{Q}f | f \rangle + \langle \hat{Q}g | g \rangle + \langle \hat{Q}f | g \rangle + \langle \hat{Q}g | f \rangle$$

Using the assumption  $\langle h | \hat{Q}h \rangle = \langle \hat{Q}h | h \rangle$  for all functions, we have  $\langle f | \hat{Q}f \rangle = \langle \hat{Q}f | f \rangle$  and  $\langle g | \hat{Q}g \rangle = \langle \hat{Q}g | g \rangle$  so, equating the two expansions above and cancelling terms, we get

$$(0.8) \quad \langle f|\hat{Q}g\rangle + \langle g|\hat{Q}f\rangle = \langle \hat{Q}f|g\rangle + \langle \hat{Q}g|f\rangle$$

Repeating this procedure for  $h = f + ig$ , we have:

$$(0.9) \quad \langle h|\hat{Q}h\rangle = \langle f + ig|\hat{Q}f + i\hat{Q}g\rangle$$

$$(0.10) \quad = \langle f|\hat{Q}f\rangle + (-i)(i)\langle g|\hat{Q}g\rangle + i\langle f|\hat{Q}g\rangle - i\langle g|\hat{Q}f\rangle$$

$$(0.11) \quad = \langle f|\hat{Q}f\rangle + \langle g|\hat{Q}g\rangle + i\langle f|\hat{Q}g\rangle - i\langle g|\hat{Q}f\rangle$$

Here, the factor of  $-i$  in the second term comes from the complex conjugate of  $ig$ .

Finally, we calculate  $\langle \hat{Q}h|h\rangle$  for  $h = f + ig$  to get

$$(0.12) \quad \langle \hat{Q}h|h\rangle = \langle \hat{Q}f|f\rangle + \langle \hat{Q}g|g\rangle + i\langle \hat{Q}f|g\rangle - i\langle \hat{Q}g|f\rangle$$

(The  $+i$  and  $-i$  multiply to give 1 in the second term  $\langle \hat{Q}g|g\rangle$ .)

Equating the last two expansions, as before, leads to:

$$(0.13) \quad \langle f|\hat{Q}g\rangle - \langle g|\hat{Q}f\rangle = \langle \hat{Q}f|g\rangle - \langle \hat{Q}g|f\rangle$$

If we now add equations 0.8 and 0.13 we get:

$$(0.14) \quad \langle f|\hat{Q}g\rangle = \langle \hat{Q}f|g\rangle$$

Similarly if we subtract 0.13 from 0.8 we get:

$$(0.15) \quad \langle g|\hat{Q}f\rangle = \langle \hat{Q}g|f\rangle$$

Thus we reclaim the more general Hermitian condition we started with.

#### PINGBACKS

Pingback: Hermitian operators - a few theorems

Pingback: Anti-hermitian operators