

HERMITIAN OPERATORS: PERIODIC FUNCTION

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Post date: 10 Sep 2012.

References: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Section 3.2.2; Problem 3.8.

As a simple example of the fact that, for a hermitian operator, the eigenvalues are real and, for distinct eigenvalues, the eigenfunctions are orthogonal, we'll consider again the periodic function we looked at earlier.

For the operator $d^2/d\phi^2$, the eigenvalues are the negatives of the squares of the integers, so they are obviously real. The eigenfunctions are

$$f = e^{in\phi} \quad (1)$$

where n is any integer (positive, negative or zero).

To show they are orthogonal, we calculate

$$\int_0^{2\pi} e^{in_1\phi} e^{-in_2\phi} d\phi = \int_0^{2\pi} e^{i(n_1-n_2)\phi} d\phi \quad (2)$$

$$= \frac{1}{i(n_1-n_2)} e^{i(n_1-n_2)\phi} \Big|_0^{2\pi} \quad (3)$$

$$= 0 \quad (4)$$

provided $n_1 \neq n_2$.

Remember that the eigenvalues of this operator are actually $(in)^2 = -n^2$, so in this case the two eigenfunctions (one for in and the other for $-in$) for the same eigenvalue also happen to be orthogonal.

If we look at the operator $id/d\phi$, we find the eigenvalues are all integers, with eigenfunctions $e^{in\phi}$ since

$$i \frac{d}{d\phi} e^{in\phi} = -ne^{in\phi} \quad (5)$$

This time, the eigenvalues are all non-degenerate. The eigenfunctions are all mutually orthogonal by the same argument as above.

PINGBACKS

Pingback: Momentum: eigenvalues and normalization