

## MOMENTUM: EIGENVALUES AND NORMALIZATION

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Section 3.3.2; Problem 3.9.

The example of a periodic function which we studied earlier had discrete eigenvalues for both the first and second derivative of the periodic variable. In particular, for the operator  $i\hbar d/d\phi$  we found that the eigenvalues are all integers, with eigenfunctions  $e^{in\phi}$  since

$$(1) \quad i \frac{d}{d\phi} e^{in\phi} = -ne^{in\phi}$$

This operator bears a strong resemblance to the momentum operator in one dimension, which is  $\hat{p} = -i\hbar d/dx$ . However, if we try to find the eigenvalues and eigenfunctions of  $\hat{p}$ , we run into a bit of a problem. We try to solve, for some eigenvalue  $p$ :

$$(2) \quad \hat{p}f = pf$$

$$(3) \quad -i\hbar \frac{d}{dx} f = pf$$

This has the solution

$$(4) \quad f_p(x) = Ae^{ipx/\hbar}$$

for some constant  $A$ . Ordinarily, at this stage, we would impose some boundary condition on the solution to obtain acceptable values of  $p$ . The problem is that we'd like to define this function over all  $x$  and, if we try to do this, the function is not normalizable for any value of  $p$ . At first glance, we might think that if we chose  $p$  to be purely imaginary as in  $p = \alpha i$ , it might work since we get

$$(5) \quad f(x) = Ae^{-\alpha x/\hbar}$$

but of course this tends to infinity at large negative  $x$  so that doesn't work.

In fact if  $p$  has a non-zero imaginary part,  $f(x)$  goes to infinity at one end of its domain. So we're restricted to looking at real values of  $p$ .

In that case,  $f(x)$  is periodic and thus is still not normalizable. Thus there are no eigenfunctions of the momentum operator that lie in Hilbert space (which, remember, is the vector space of square-integrable functions).

What happens if do the normalization integral anyway? That is, we try

$$(6) \quad \int_{-\infty}^{\infty} f_{p_1}^*(x) f_{p_2}(x) dx = |A|^2 \int_{-\infty}^{\infty} e^{i(p_2-p_1)x/\hbar} dx$$

By using the variable transformation  $\xi \equiv x/\hbar$ , we get

$$(7) \quad \int_{-\infty}^{\infty} f_{p_1}^*(x) f_{p_2}(x) dx = |A|^2 \hbar \int_{-\infty}^{\infty} e^{i(p_2-p_1)\xi} d\xi$$

It's at this point that we invoke the dodgy formula involving the Dirac delta function that we obtained a while back. Using this, we can write the integral as a delta function, and we get

$$(8) \quad \int_{-\infty}^{\infty} f_{p_1}^*(x) f_{p_2}(x) dx = 2\pi |A|^2 \hbar \delta(p_2 - p_1)$$

This is sort of like a normalization condition, in that the integral is zero when  $p_1 \neq p_2$  (that is, if you believe that the integral really does evaluate to a delta function), and non-zero (infinite, in fact) if  $p_1 = p_2$ . In fact, if we take the constant  $A$  to be

$$(9) \quad A = \frac{1}{\sqrt{2\pi\hbar}}$$

and use the bra-ket notation for the integral, we can write

$$(10) \quad \langle f_{p_1} | f_{p_2} \rangle = \delta(p_2 - p_1)$$

We can also express an arbitrary function  $g(x)$  as a Fourier transform over  $p$  by writing

$$(11) \quad g(x) = \int_{-\infty}^{\infty} c(p) f_p(x) dp$$

$$(12) \quad = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} c(p) e^{ipx/\hbar} dp$$

$$(13) \quad g(\hbar\xi) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} c(p) e^{ip\xi} dp$$

From Plancherel's theorem, we can invert this relation to get  $c(p)$ :

$$(14) \quad c(p) = \sqrt{\frac{\hbar}{2\pi}} \int_{-\infty}^{\infty} g(\hbar\xi) e^{-ip\xi} d\xi$$

$$(15) \quad = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} g(x) e^{-ipx/\hbar} dx$$

$$(16) \quad = \langle f_p | g \rangle$$

In general, hermitian operators with continuous eigenvalues don't have normalizable eigenfunctions and have to be analyzed in this way. In particular, the hamiltonian (energy) of a system can have an entirely discrete spectrum (infinite square well or harmonic oscillator), a totally continuous spectrum (free particle, delta function barrier or finite square barrier) or a mixture of the two (delta function well or finite square well).

#### PINGBACKS

Pingback: Infinite square well: momentum

Pingback: Momentum space: harmonic oscillator

Pingback: Momentum space: mean position

Pingback: Sequential measurements

Pingback: Momentum space representation of finite wave function

Pingback: Momentum space: another example

Pingback: Free particle in momentum space

Pingback: Non-denumerable basis: position and momentum states

Pingback: Differential operators - matrix elements and hermiticity

Pingback: Postulates of quantum mechanics: momentum

Pingback: Parity transformations

Pingback: Position operator - eigenfunctions

Pingback: Eigenfunctions of position and momentum; unit operators