

## COMMUTATORS: A FEW THEOREMS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 3.13.

The commutator of two operators is defined as

$$(0.1) \quad [A, B] \equiv AB - BA$$

In general, a commutator is non-zero, since the order in which we apply operators can make a difference. In practice, to work out a commutator we need to apply it to a test function  $f$ , so that we really need to work out  $[A, B]f$  and then remove the test function to see the result. This is because many operators, such as the momentum, involve taking the derivative.

We'll now have a look at a few theorems involving commutators.

Theorem 1:

$$(0.2) \quad [AB, C] = A[B, C] + [A, C]B$$

Proof: The LHS is:

$$(0.3) \quad [AB, C] = ABC - CAB$$

The RHS is:

$$(0.4) \quad A[B, C] + [A, C]B = ABC - ACB + ACB - CAB$$

$$(0.5) \quad = ABC - CAB$$

$$(0.6) \quad = [AB, C]$$

QED.

Theorem 2:

$$(0.7) \quad [x^n, p] = i\hbar nx^{n-1}$$

where  $p$  is the momentum operator.

Proof: Using  $p = \frac{\hbar}{i}\partial/\partial x$  and letting the commutator operate on some arbitrary function  $g$ :

$$(0.8) \quad [x^n, p]g = x^n \frac{\hbar}{i} \frac{\partial g}{\partial x} - \frac{\hbar}{i} \frac{\partial}{\partial x}(x^n g)$$

$$(0.9) \quad = x^n \frac{\hbar}{i} \frac{\partial g}{\partial x} - \frac{\hbar}{i} n x^{n-1} g - x^n \frac{\hbar}{i} \frac{\partial g}{\partial x}$$

$$(0.10) \quad = i\hbar n x^{n-1} g$$

Removing the function  $g$  gives the result  $[x^n, p] = i\hbar n x^{n-1}$ . QED.

Theorem 3:

$$(0.11) \quad [f(x), p] = i\hbar \frac{df}{dx}$$

Again, letting the commutator operate on a function  $g$ :

$$(0.12) \quad [f(x), p] = f \frac{\hbar}{i} \frac{\partial g}{\partial x} - \frac{\hbar}{i} \frac{\partial}{\partial x}(fg)$$

$$(0.13) \quad = f \frac{\hbar}{i} \frac{\partial g}{\partial x} - \frac{\hbar}{i} \frac{\partial f}{\partial x} g - f \frac{\hbar}{i} \frac{\partial g}{\partial x}$$

$$(0.14) \quad = i\hbar \frac{\partial f}{\partial x} g$$

Removing  $g$  gives the result  $[f(x), p] = i\hbar \partial f / \partial x$ . QED.

#### PINGBACKS

Pingback: Angular momentum - commutators with position and momentum

Pingback: Selection rules for spontaneous emission of radiation

Pingback: Lie brackets (commutators)

Pingback: The classical limit of quantum mechanics; Ehrenfest's theorem

Pingback: Poisson brackets to commutators: classical to quantum

Pingback: Translational invariance and conservation of momentum

Pingback: Finite transformations: correspondence between classical and quantum

Pingback: Parity transformations