

UNCERTAINTY PRINCIPLE - EXAMPLES

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Post date: 25 Jul 2011.

Reference: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Sec 3.5; Problem 3.14.

In this post we'll have a look at an example involving the uncertainty principle. This makes use of the generic uncertainty principle for two observables A and B :

$$\sigma_A^2 \sigma_B^2 \geq \left(\frac{1}{2i} \langle [\hat{A}, \hat{B}] \rangle \right)^2 \quad (1)$$

We can calculate an expression for a 'position-energy' uncertainty relation. For this we'll need the commutator $[\hat{x}, \hat{H}]$, so we work that out first. Since H involves a derivative, we use a test function g on which to operate:

$$[\hat{x}, \hat{H}]g = -\frac{\hbar^2}{2m}x \frac{\partial^2 g}{\partial x^2} + xVg + -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2}(xg) - xVg \quad (2)$$

$$= \frac{\hbar^2}{2m} \left(-x \frac{\partial^2 g}{\partial x^2} + 2 \frac{\partial g}{\partial x} + x \frac{\partial^2 g}{\partial x^2} \right) \quad (3)$$

$$= \frac{\hbar^2}{m} \frac{\partial g}{\partial x} \quad (4)$$

$$= \frac{i\hbar}{m} (pg) \quad (5)$$

From 1 with $\hat{A} = \hat{x}$ and $\hat{B} = \hat{H}$ we have

$$\sigma_x^2 \sigma_H^2 \geq \left(\frac{1}{2i} \langle [\hat{x}, \hat{H}] \rangle \right)^2 \quad (6)$$

$$= \frac{\hbar^2}{4m^2} \langle p \rangle^2 \quad (7)$$

so the uncertainty principle here becomes

$$\sigma_x \sigma_H \geq \frac{\hbar}{2m} |\langle p \rangle| \quad (8)$$

For a stationary state, this doesn't tell you much because the average position of the particle doesn't change, so $\langle p \rangle = 0$. For linear combinations of stationary states, though, the average momentum will not be zero, so this condition is more than trivial.

PINGBACKS

Pingback: [Uncertainty principle: rates of change of operators](#)

Pingback: [Energy-time uncertainty principle - example](#)