

MATRIX ELEMENTS: EXAMPLE

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

Post date: 30 Sep 2012.

References: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Section 3.6; Problem 3.22.

As a simple example of the bra-ket notation and the matrix representation of operators, suppose we have a three-dimensional vector space spanned by an orthonormal basis $|1\rangle, |2\rangle, |3\rangle$. We begin with a couple of vectors given by

$$|\alpha\rangle = i|1\rangle - 2|2\rangle - i|3\rangle \quad (1)$$

$$|\beta\rangle = i|1\rangle + 2|3\rangle \quad (2)$$

The corresponding bras are found by taking the complex conjugate:

$$\langle\alpha| = -i\langle 1| - 2\langle 2| + i\langle 3| \quad (3)$$

$$\langle\beta| = -i\langle 1| + 2\langle 3| \quad (4)$$

Remember that these bras are not vectors in their own right; rather, they are operators which acquire meaning only when they are applied to vectors to yield a complex number.

We can form the two inner products of these two vectors:

$$\langle\alpha|\beta\rangle = (-i)(i) + (-2)(0) + (i)(2) \quad (5)$$

$$= 1 + 2i \quad (6)$$

$$\langle\beta|\alpha\rangle = (-i)(i) + (0)(-2) + (2)(-i) \quad (7)$$

$$= 1 - 2i \quad (8)$$

Clearly $\langle\beta|\alpha\rangle = \langle\alpha|\beta\rangle^*$.

If we define an operator $\hat{A} = |\alpha\rangle\langle\beta|$ we can write it in terms of the basis above:

$$\hat{A} = |\alpha\rangle\langle\beta| \quad (9)$$

$$= (i|1\rangle - 2|2\rangle - i|3\rangle)(-i\langle 1| + 2\langle 3|) \quad (10)$$

$$= |1\rangle\langle 1| + 2i|1\rangle\langle 3| + 2i|2\rangle\langle 1| - 4|2\rangle\langle 3| - |3\rangle\langle 1| - 2i|3\rangle\langle 3| \quad (11)$$

From here, we can obtain its matrix elements in this basis by using the orthonormal property of the basis. For example

$$\langle 1|A|1\rangle = \langle 1|1\rangle\langle 1|1\rangle + 2i\langle 1|1\rangle\langle 3|1\rangle + 2i\langle 1|2\rangle\langle 1|1\rangle \quad (12)$$

$$- 4\langle 1|2\rangle\langle 3|1\rangle - \langle 1|3\rangle\langle 1|1\rangle - 2i\langle 1|3\rangle\langle 3|1\rangle \quad (13)$$

$$= 1 \quad (14)$$

Doing similar calculations for the other elements, we get

$$A = \begin{bmatrix} 1 & 0 & 2i \\ 2i & 0 & -4 \\ -1 & 0 & -2i \end{bmatrix} \quad (15)$$

The matrix is not hermitian, as it is not equal to its conjugate transpose.