

RIGID ROTOR IN QUANTUM MECHANICS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education Problem 4.24.

A rigid rotor is a system in which we have two particles, each of mass m , at the ends of a rigid rod of length a . The centre of the rod is fixed, but the rod is free to rotate about its centre in any direction (so it's not rotating on a fixed axis). Assuming no potential energy, the energy of the system is, for the two masses

$$(0.1) \quad E = 2 \frac{p^2}{2m} = \frac{p^2}{m}$$

Since the particles are constrained to have rotational motion only, and the rod is fixed, \mathbf{r} is always perpendicular to \mathbf{p} , so taking \mathbf{p} to be the momentum of one of the masses, $|\mathbf{L}| = 2|\mathbf{r}||\mathbf{p}| = 2(a/2)p = ap$ and $L^2 = a^2p^2$ so

$$(0.2) \quad E = \frac{L^2}{ma^2}$$

We know the eigenvalues of L^2 are $\hbar^2 n(n+1)$ for $n = 0, 1, 2, \dots$ so

$$(0.3) \quad E_n = \frac{\hbar^2 n(n+1)}{ma^2}$$

Since E is directly proportional to L^2 , the eigenfunctions are just the spherical harmonics, and state n therefore has degeneracy $2n + 1$.