

SPIN: THE X AND Y COMPONENTS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

Reference: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 4.29.

Just as we found the eigenvalues and eigenspinors of S_z , we can do the same for the other 2 components. For S_y , for example, we have

$$S_y = \frac{\hbar}{2} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \quad (1)$$

We have, first, for the eigenvalues

$$\begin{vmatrix} -\lambda & -i\frac{\hbar}{2} \\ i\frac{\hbar}{2} & -\lambda \end{vmatrix} = \lambda^2 - \frac{\hbar^2}{4} = 0 \quad (2)$$

from which

$$\lambda = \pm \frac{\hbar}{2} \quad (3)$$

So the eigenvalues of S_y are also $\pm\hbar/2$.

Solving the equation to find the eigenspinors gives

$$\frac{\hbar}{2} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \pm \frac{\hbar}{2} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} \quad (4)$$

$$\chi_+^{(y)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \end{pmatrix} \quad (5)$$

$$\chi_-^{(y)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \end{pmatrix} \quad (6)$$

A similar calculation for S_x gives

$$\chi_+^{(x)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (7)$$

$$\chi_-^{(x)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix} \quad (8)$$

To find the probabilities of a particle that is in a general spin state $\chi = \begin{pmatrix} a \\ b \end{pmatrix}$ being in each of the eigenstates, we need to express the general state χ in terms of $\chi_+^{(y)}$ and $\chi_-^{(y)}$. First we convert the basis vectors:

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} = \frac{1}{\sqrt{2}}(\chi_+^{(y)} + \chi_-^{(y)}) \quad (9)$$

$$\begin{pmatrix} 0 \\ 1 \end{pmatrix} = \frac{-i}{\sqrt{2}}(\chi_+^{(y)} - \chi_-^{(y)}) \quad (10)$$

so

$$\chi = a \begin{pmatrix} 1 \\ 0 \end{pmatrix} + b \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \frac{1}{\sqrt{2}}(a - ib)\chi_+^{(y)} + \frac{1}{\sqrt{2}}(a + ib)\chi_-^{(y)} \quad (11)$$

Thus the probability of the particle being in state $\chi_+^{(y)}$ is $|a - ib|^2/2$ and in state $\chi_-^{(y)}$ is $|a + ib|^2/2$.

The sum of these probabilities is (remember that a and b are, in general, complex, so we can't just say $|a + ib|^2 = a^2 + b^2$)

$$\frac{|a - ib|^2}{2} + \frac{|a + ib|^2}{2} = \frac{1}{2}(|a|^2 + |b|^2 - ia^*b + iab^* + |a|^2 + |b|^2 + ia^*b - iab^*) \quad (12)$$

$$= \frac{1}{2}(2|a|^2 + 2|b|^2) \quad (13)$$

$$= 1 \quad (14)$$

since $|a|^2 + |b|^2 = 1$ from normalization.

Since $S_x^2 = S_y^2 = S_z^2 = \frac{\hbar^2}{4} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, measuring S_y^2 will always give $\hbar^2/4$.

Since the eigenvalues of S_y are $\pm\hbar/2$ this follows.

PINGBACKS

Pingback: Spin 1/2 particle in a magnetic field

Pingback: Spin 1/2: spin components

Pingback: Stern-gerlach experiment