

SPIN 1/2: SPIN COMPONENTS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

Reference: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 4.49.

As an example of some calculations with a spin 1/2 system, suppose we have an electron in a spin state given by

$$\chi = A \begin{pmatrix} 1-2i \\ 2 \end{pmatrix} \quad (1)$$

Requiring $|\chi|^2 = 1$ gives $|A|^2(5+4) = 9|A|^2 = 1$, so if we take A as real, then $A = 1/3$.

Now suppose we measured each spin component on this system; what values could we get? Note that we're considering measuring a component, then restoring the system to its initial state, and then measuring the next component. If we didn't do a restoration after each measurement, we would alter the state to the eigenstate of the measured value, which we don't want to do.

First, we'll look at S_z . We need to express χ as a linear combination of the eigenspinors of S_z so:

$$\chi = a \begin{pmatrix} 1 \\ 0 \end{pmatrix} + b \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (2)$$

Comparison with the original state gives:

$$\chi = \frac{1-2i}{3} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \frac{2}{3} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (3)$$

so the possible values are $\hbar/2$ with probability $|(1-2i)/3|^2 = 5/9$ and $-\hbar/2$ with probability $4/9$.

To find the expectation value $\langle S_z \rangle$ we need to work out $\chi^T S_z \chi$.

$$\frac{1}{9} \begin{pmatrix} 1+2i & 2 \end{pmatrix} \frac{\hbar}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1-2i \\ 2 \end{pmatrix} = \frac{\hbar}{18} (5-4) = \frac{\hbar}{18} \quad (4)$$

For S_x we use the eigenspinors from our earlier post

$$\chi = \frac{\sqrt{2}}{3} \left[a \begin{pmatrix} 1 \\ 1 \end{pmatrix} + b \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right] \quad (5)$$

Solving for a and b :

$$a + b = 1 - 2i \quad (6)$$

$$a - b = 2 \quad (7)$$

$$a = \frac{3}{2} - i \quad (8)$$

$$b = -\frac{1}{2} - i \quad (9)$$

so

$$\chi = \frac{\sqrt{2}}{3} \left[\left(\frac{3}{2} - i \right) \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \left(-\frac{1}{2} - i \right) \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right] \quad (10)$$

The possible values are $\hbar/2$ with probability $|(\sqrt{2}/3)(3/2 - i)|^2 = 13/18$ and $-\hbar/2$ with probability $5/18$.

The expectation value is found from the matrix $S_x = \frac{\hbar}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

$$\langle S_x \rangle = \frac{1}{9} \begin{pmatrix} 1 + 2i & 2 \end{pmatrix} \frac{\hbar}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 - 2i \\ 2 \end{pmatrix} = \frac{\hbar}{18} (2 + 4i + 2 - 4i) = \frac{2\hbar}{9} \quad (11)$$

For S_y we can use the eigenspinors from before, and following the same procedure as above we get

$$\chi = \frac{\sqrt{2}}{3} \left[\left(\frac{1}{2} - 2i \right) \begin{pmatrix} 1 \\ i \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 1 \\ -i \end{pmatrix} \right] \quad (12)$$

The possible values are $\hbar/2$ with probability $|(\sqrt{2}/3)(1/2 - 2i)|^2 = 17/18$ and $-\hbar/2$ with probability $1/18$. The expectation value is

$$\langle S_y \rangle = \frac{1}{9} \begin{pmatrix} 1 + 2i & 2 \end{pmatrix} \frac{\hbar}{2} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{pmatrix} 1 - 2i \\ 2 \end{pmatrix} = \frac{\hbar}{18} (-2i + 4 + 2i + 4) = \frac{4\hbar}{9} \quad (13)$$

As a check, $\langle S_x \rangle^2 + \langle S_y \rangle^2 + \langle S_z \rangle^2 = (\hbar/2)^2$. Note that this isn't the same as $\langle S_x^2 \rangle + \langle S_y^2 \rangle + \langle S_z^2 \rangle = \langle S^2 \rangle = s(s+1)\hbar^2 = \frac{3\hbar^2}{4}$.

PINGBACKS

Pingback: Angular momentum: adding spins in arbitrary directions