

HARMONIC OSCILLATOR: FIRST ORDER PERTURBATION

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 6.2.

This is a simple example of applying first order perturbation theory to the harmonic oscillator. The energy levels of an unperturbed oscillator are

$$E_{n0} = \left(n + \frac{1}{2}\right) \hbar\omega \quad (1)$$

where $\omega = \sqrt{k/m}$ and the potential is $V = \frac{1}{2}kx^2$. If we perturb the potential by changing k slightly, so the new potential is

$$V' = \frac{1}{2}(1 + \epsilon)kx^2 \quad (2)$$

then, of course, it's easy to find the exact energy levels just by changing k in the original formula:

$$E_n = \left(n + \frac{1}{2}\right) \hbar\omega' \quad (3)$$

$$= \left(n + \frac{1}{2}\right) \hbar\sqrt{\frac{k(1 + \epsilon)}{m}} \quad (4)$$

We can expand the square root in a power series:

$$E_n = \left(n + \frac{1}{2}\right) \hbar\sqrt{\frac{k}{m}} \left[1 + \frac{\epsilon}{2} - \frac{\epsilon^2}{8} + \dots\right] \quad (5)$$

From first order perturbation theory, the change to the energy is (since the perturbation in the potential here is $V' - V = \frac{1}{2}\epsilon kx^2 = \epsilon V$):

$$E_{n1} = \langle n0 | \epsilon V | n0 \rangle \quad (6)$$

We could do the integral implied here, but we've already worked out the mean value of the potential for the harmonic oscillator using the virial theorem, and we know that $\langle V \rangle = \langle T \rangle = E_n/2$ so

$$E_{n1} = \frac{\epsilon}{2} E_{n0} \quad (7)$$

$$= \frac{\epsilon}{2} \left(n + \frac{1}{2} \right) \hbar \sqrt{\frac{k}{m}} \quad (8)$$

This is the first order term in ϵ in the series expansion above.

PINGBACKS

Pingback: Second order non-degenerate perturbation theory