

## ZEEMAN EFFECT: THE N=2 LINE IN HYDROGEN

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

References: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 6.23.

We'll show how the strong field Zeeman effect alters the spectrum of hydrogen for the case  $n = 2$ . The full formula for the energy levels is

$$E_n = -\frac{13.6 \text{ eV}}{n^2} \left[ 1 - \frac{\alpha^2}{n} \left( \frac{3}{4n} - \frac{\ell(\ell+1) - \ell_z s_z}{\ell(\ell+\frac{1}{2})(\ell+1)} \right) \right] + \mu_B (\ell_z + 2s_z) B_{ext} \quad (1)$$
$$= E_{2,0} + E_{fs1} + E_{Z1} \quad (2)$$

which is the sum of the Bohr energy, and the fine structure and the Zeeman corrections.

For  $n = 2$ , the Bohr energy is

$$E_{2,0} = -\frac{13.6 \text{ eV}}{2^2} = -3.4 \text{ eV} \quad (3)$$

and this is the same for all substates.

The fine structure term is

$$E_{fs1} = (3.4 \text{ eV}) \frac{\alpha^2}{2} \left( \frac{3}{8} - \frac{\ell(\ell+1) - \ell_z s_z}{\ell(\ell+\frac{1}{2})(\ell+1)} \right) \quad (4)$$

If  $\ell = 0$ , the second term in the parentheses is zero divided by zero (since the only possible value of  $\ell_z$  is zero in this case), but as we'll see in the next post, we can take this quotient to be 1. For  $\ell = 1$  we can just plug in the numbers and get

$$E_{fs1}(\ell = 1) = (3.4 \text{ eV}) \frac{\alpha^2}{2} \left( \frac{3}{8} - \frac{2 - \ell_z s_z}{3} \right) \quad (5)$$

. The Zeeman term depends only on the  $z$  components. The results are

$\ell$	$\ell_z$	$s_z$	$E_{Z1} (\times \mu_B B_{ext})$	$E_{fs1} (\times 3.4\alpha^2, \text{ eV})$	$E_n$
0	0	$\frac{1}{2}$	1	$-\frac{5}{16} = -0.3125$	$-3.4 \left(1 + \frac{5}{16}\alpha^2\right) + \mu_B B_{ext}$
0	0	$-\frac{1}{2}$	-1	$-\frac{5}{16} = -0.3125$	$-3.4 \left(1 + \frac{5}{16}\alpha^2\right) - \mu_B B_{ext}$
1	-1	$\frac{1}{2}$	0	$-\frac{11}{48} = -0.2292$	$-3.4 \left(1 + \frac{11}{48}\alpha^2\right)$
1	-1	$-\frac{1}{2}$	-2	$-\frac{1}{16} = -0.0625$	$-3.4 \left(1 + \frac{1}{16}\alpha^2\right) - 2\mu_B B_{ext}$
1	0	$\frac{1}{2}$	1	$-\frac{7}{48} = -0.1458$	$-3.4 \left(1 + \frac{7}{48}\alpha^2\right) + \mu_B B_{ext}$
1	0	$-\frac{1}{2}$	-1	$-\frac{7}{48} = -0.1458$	$-3.4 \left(1 + \frac{7}{48}\alpha^2\right) - \mu_B B_{ext}$
1	1	$\frac{1}{2}$	2	$-\frac{1}{16} = -0.0625$	$-3.4 \left(1 + \frac{1}{16}\alpha^2\right) + 2\mu_B B_{ext}$
1	1	$-\frac{1}{2}$	0	$-\frac{11}{48} = -0.2292$	$-3.4 \left(1 + \frac{11}{48}\alpha^2\right)$

Looking only at the Zeeman energies, there are 5 distinct energies, 3 of which have degeneracy 2 and the other 2 of which have degeneracy 1. Two of the states are unaffected by the external magnetic field.

## PINGBACKS

Pingback: Zeeman effect: degenerate perturbation theory for  $n = 2$