

## WKB APPROXIMATION - ALTERNATIVE DERIVATION

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

References: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 8.2.

Here's an alternative derivation of the WKB approximation. This time we start with the wave function in the form

$$(0.1) \quad \psi(x) = e^{if(x)/\hbar}$$

where  $f(x)$  is now a complex function. Note that we don't need a separate amplitude term multiplying the exponential, since the imaginary part of  $f$  gives a real function:

$$(0.2) \quad e^{if(x)/\hbar} = e^{-Im(f(x))/\hbar} e^{iRe(f(x))/\hbar}$$

We can plug this into the one-dimensional time-independent Schrödinger equation

$$(0.3) \quad -\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} + V(x)\psi = E\psi$$

and rewrite it as

$$(0.4) \quad \frac{d^2\psi}{dx^2} = -\frac{2m[E - V(x)]}{\hbar^2}\psi$$

$$(0.5) \quad \equiv -\frac{p^2}{\hbar^2}\psi$$

where  $p = \sqrt{2m[E - V(x)]}$  is the classical formula for the momentum of a particle with total energy  $E$  moving in a one-dimensional potential  $V(x)$ , provided we assume  $E \geq V(x)$  for all  $x$ . The derivatives are

$$(0.6) \quad \psi' = e^{if/\hbar} \frac{if'}{\hbar}$$

$$(0.7) \quad = \frac{if'}{\hbar} \psi$$

$$(0.8) \quad \psi'' = \frac{if''}{\hbar} \psi + \frac{if'}{\hbar} \psi'$$

$$(0.9) \quad = \frac{if''}{\hbar} \psi - \left(\frac{f'}{\hbar}\right)^2 \psi$$

Putting this into 0.5 we get

$$(0.10) \quad \frac{if''}{\hbar} \psi - \left(\frac{f'}{\hbar}\right)^2 \psi = -\frac{p^2}{\hbar^2} \psi$$

$$(0.11) \quad i\hbar f'' - (f')^2 + p^2 = 0$$

Since  $p$  can be an arbitrary function of  $x$ , we can't usually solve this differential equation in closed form, but we can try a series solution. If we expand  $f(x)$  in powers of  $\hbar$  we have

$$(0.12) \quad f(x) = \sum_{i=0}^{\infty} \hbar^i f_i(x)$$

so we can substitute this into 0.11 and require the coefficient of each power of  $\hbar$  to be zero:

$$(0.13) \quad i \sum_{i=1}^{\infty} \hbar^i f''_{i-1} - \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \hbar^{i+j} f'_i f'_j + p^2 = 0$$

$$(0.14) \quad -(f'_0)^2 + p^2 = 0$$

$$(0.15) \quad if''_0 - 2f'_0 f'_1 = 0$$

$$(0.16) \quad if''_1 - 2f'_0 f'_2 - (f'_1)^2 = 0$$

Equation 0.14 can be solved as

$$(0.17) \quad f_0(x) = \pm \int p(x) dx$$

$$(0.18) \quad f''_0 = \pm p'(x)$$

Then from 0.15 we get

$$(0.19) \quad \pm ip' = \pm 2pf_1'$$

$$(0.20) \quad \frac{ip'}{2p} = f_1'$$

$$(0.21) \quad \frac{i}{2} \ln p = f_1 + \ln C$$

$$(0.22) \quad if_1 = -\frac{1}{2} \ln p + \ln C$$

To first order in  $\hbar$  we then get for the wave function:

$$(0.23) \quad \psi(x) = e^{i(f_0 + \hbar f_1)/\hbar}$$

$$(0.24) \quad = e^{if_1} e^{if_0/\hbar}$$

$$(0.25) \quad = \frac{C}{\sqrt{p}} e^{\pm \frac{i}{\hbar} \int p(x) dx}$$

which is the same formula for the WKB approximation that we got before.