

GEOMETRIC PHASE IS ALWAYS ZERO FOR REAL WAVE FUNCTIONS

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: Griffiths, David J. (2005), Introduction to Quantum Mechanics, 2nd Edition; Pearson Education - Problem 10.5.

Here's another example of calculating phases in the adiabatic theorem which says that if a system starts out in the n th state of a time-dependent hamiltonian, and the hamiltonian changes slowly compared to the internal period of the time-independent wave function (that is, the time scale over which the hamiltonian changes is much longer than \hbar/E_n), then after a time t the system will end up in state

$$\Psi_n(t) = e^{i\theta_n(t)} e^{i\gamma_n(t)} \psi_n(t) \quad (1)$$

where

$$\theta_n(t) \equiv -\frac{1}{\hbar} \int_0^t E_n(t') dt' \quad (2)$$

$$\gamma_n(t) \equiv i \int_0^t \left\langle \psi_n(t') \left| \frac{\partial}{\partial t'} \psi_n(t') \right. \right\rangle dt' \quad (3)$$

θ is called the *dynamic phase* and γ is called the *geometric phase*. If ψ_n is real, then γ_n is always zero, as we can see by differentiating the normalization condition:

$$\langle \psi_n | \psi_n \rangle = 1 \quad (4)$$

$$\frac{d}{dt} \langle \psi_n | \psi_n \rangle = 0 \quad (5)$$

$$= \langle \dot{\psi}_n | \psi_n \rangle + \langle \psi_n | \dot{\psi}_n \rangle \quad (6)$$

$$= \langle \psi_n | \dot{\psi}_n \rangle^* + \langle \psi_n | \dot{\psi}_n \rangle \quad (7)$$

$$= 2\Re(\langle \psi_n | \dot{\psi}_n \rangle) \quad (8)$$

That is, $\langle \psi_n(t') | \frac{\partial}{\partial t'} \psi_n(t') \rangle$ must be purely imaginary, so if ψ_n is real, the bracket must be zero. This also means that γ is always real.

We can multiply the real wave function ψ_n by a phase factor $e^{i\phi_n}$ where ϕ_n is a real function of whatever parameters are dependent on time in the hamiltonian (but ϕ_n is not a function of x). In that case we have a new wave function (we'll drop the subscript n to save time):

$$\psi' = e^{i\phi}\psi \quad (9)$$

$$\dot{\psi}' = i\dot{\phi}e^{i\phi}\psi + e^{i\phi}\dot{\psi} \quad (10)$$

$$\langle \psi' | \dot{\psi}' \rangle = \langle \psi | i\dot{\phi}\psi + \dot{\psi} \rangle \quad (11)$$

$$= i\langle \psi | \dot{\phi}\psi \rangle + \langle \psi | \dot{\psi} \rangle \quad (12)$$

$$= i\dot{\phi} \quad (13)$$

where in the last line we took $\dot{\phi}$ outside the bracket since it doesn't depend on x and used $\langle \psi | \dot{\psi} \rangle = 0$. The geometric phase for the modified wave function is therefore

$$\gamma' = i \int_0^t i\dot{\phi} dt' \quad (14)$$

$$= -(\phi(t) - \phi(0)) \quad (15)$$

Putting this back into 1 we get

$$\Psi'(t) = e^{i\theta(t)} e^{-i(\phi(t) - \phi(0))} \psi'(t) \quad (16)$$

$$= e^{i\theta(t)} e^{-i(\phi(t) - \phi(0))} e^{i\phi(t)} \psi(t) \quad (17)$$

$$= e^{i\theta(t)} e^{i\phi(0)} \psi(t) \quad (18)$$

Although the wave function picks up a constant phase $\phi(0)$, there is no time-dependent geometric phase.

PINGBACKS

Pingback: Forced harmonic oscillator: exact solution and adiabatic approximation