

CREATION AND DESTRUCTION OPERATORS FOR A FREE SCALAR FIELD

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: References: Robert D. Klauber, *Student Friendly Quantum Field Theory*, (Sandtrove Press, 2013) - Chapter 3, Problem 3.10.

The scalar, free-field Hamiltonian derived earlier is

$$H = \sum_{\mathbf{k}} \omega_{\mathbf{k}} \left[N_a(\mathbf{k}) + \frac{1}{2} + N_b(\mathbf{k}) + \frac{1}{2} \right] \quad (1)$$

where the operators are

$$N_a(\mathbf{k}) = a^\dagger(\mathbf{k}) a(\mathbf{k}) \quad (2)$$

$$N_b(\mathbf{k}) = b^\dagger(\mathbf{k}) b(\mathbf{k}) \quad (3)$$

These operators can be interpreted as number operators. That is, when they operate on a state $|n_{\mathbf{k}}\rangle$ containing $n_{\mathbf{k}}$ particles with energy $\omega_{\mathbf{k}}$, their eigenvalues are just $n_{\mathbf{k}}$:

$$N_a(\mathbf{k}) |n_{\mathbf{k}}\rangle = n_{\mathbf{k}} |n_{\mathbf{k}}\rangle \quad (4)$$

With this interpretation, and the commutation relations for the a and b operators, we can derive the fact that $a^\dagger(\mathbf{k})$ is a *creation operator* for type a particles with energy $\omega_{\mathbf{k}}$, $a(\mathbf{k})$ is a *destruction operator* for the same type of particle, and $b^\dagger(\mathbf{k})$ and $b(\mathbf{k})$ are the analogous operators for type b particles (antiparticles).

Suppose we operate on the state $|n_{\mathbf{k}}\rangle$ with $a^\dagger(\mathbf{k})$. To see that this creates a particle with energy $\omega_{\mathbf{k}}$, we operate on the resulting state with $N_a(\mathbf{k})$:

$$N_a(\mathbf{k}) a^\dagger(\mathbf{k}) |n_{\mathbf{k}}\rangle = a^\dagger(\mathbf{k}) a(\mathbf{k}) a^\dagger(\mathbf{k}) |n_{\mathbf{k}}\rangle \quad (5)$$

$$= a^\dagger(\mathbf{k}) \left[1 + a^\dagger(\mathbf{k}) a(\mathbf{k}) \right] |n_{\mathbf{k}}\rangle \quad (6)$$

$$= a^\dagger(\mathbf{k}) [1 + N_a(\mathbf{k})] |n_{\mathbf{k}}\rangle \quad (7)$$

$$= a^\dagger(\mathbf{k}) [1 + n_{\mathbf{k}}] |n_{\mathbf{k}}\rangle \quad (8)$$

$$= [1 + n_{\mathbf{k}}] a^\dagger(\mathbf{k}) |n_{\mathbf{k}}\rangle \quad (9)$$

Thus the state $a^\dagger(\mathbf{k})|n_{\mathbf{k}}\rangle$ is an eigenstate of $N_a(\mathbf{k})$ with eigenvalue $1+n_{\mathbf{k}}$ so the operator $a^\dagger(\mathbf{k})$ has created an extra $\omega_{\mathbf{k}}$ particle.

The procedure for $a(\mathbf{k})$ is similar:

$$N_a(\mathbf{k})a(\mathbf{k})|n_{\mathbf{k}}\rangle = a^\dagger(\mathbf{k})a(\mathbf{k})a(\mathbf{k})|n_{\mathbf{k}}\rangle \quad (10)$$

$$= [a(\mathbf{k})a^\dagger(\mathbf{k}) - 1]a(\mathbf{k})|n_{\mathbf{k}}\rangle \quad (11)$$

$$= [a(\mathbf{k})N_a(\mathbf{k}) - a(\mathbf{k})]|n_{\mathbf{k}}\rangle \quad (12)$$

$$= a(\mathbf{k})[n_{\mathbf{k}} - 1]|n_{\mathbf{k}}\rangle \quad (13)$$

$$= [n_{\mathbf{k}} - 1]a(\mathbf{k})|n_{\mathbf{k}}\rangle \quad (14)$$

Thus the state $a(\mathbf{k})|n_{\mathbf{k}}\rangle$ is an eigenstate of $N_a(\mathbf{k})$ with eigenvalue $n_{\mathbf{k}} - 1$ so the operator $a(\mathbf{k})$ has destroyed a $\omega_{\mathbf{k}}$ particle.

We can represent a state containing $n_{\mathbf{k}}$ b type particles (antiparticles) by $|\bar{n}_{\mathbf{k}}\rangle$ (a bar over a symbol means it refers to antiparticles). Then

$$N_b(\mathbf{k})|\bar{n}_{\mathbf{k}}\rangle = \bar{n}_{\mathbf{k}}|\bar{n}_{\mathbf{k}}\rangle \quad (15)$$

The two derivations above are exactly the same for antiparticles, if we replace all a operators by b operators, N_a by N_b and $n_{\mathbf{k}}$ by $\bar{n}_{\mathbf{k}}$. Thus $b^\dagger(\mathbf{k})$ creates a single antiparticle with energy $\omega_{\mathbf{k}}$ and $b(\mathbf{k})$ destroys one antiparticle with that energy.

PINGBACKS

Pingback: Creation and annihilation operators: normalization

Pingback: Anticommutators, creation and annihilation operators in the Dirac equation