

CREATION AND ANNIHILATION OPERATORS: NORMALIZATION

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: References: Robert D. Klauber, *Student Friendly Quantum Field Theory*, (Sandtrove Press, 2013) - Chapter 3, Problem 3.11.

The number operators are defined in terms of the creation and annihilation operators for the free scalar Hamiltonian as

$$(1) \quad N_a(\mathbf{k}) = a^\dagger(\mathbf{k})a(\mathbf{k})$$

$$(2) \quad N_b(\mathbf{k}) = b^\dagger(\mathbf{k})b(\mathbf{k})$$

We've seen that $a^\dagger(\mathbf{k})$ creates a particle of energy $\omega_{\mathbf{k}}$ when it operates on a state, and $a(\mathbf{k})$ destroys a particle with energy $\omega_{\mathbf{k}}$ when it operates on a state (if that state contains such a particle). That is, the state $a^\dagger(\mathbf{k})|n_{\mathbf{k}}\rangle$ is an eigenstate of $N_a(\mathbf{k})$ with eigenvalue $n_{\mathbf{k}} + 1$ and $a(\mathbf{k})|n_{\mathbf{k}}\rangle$ is an eigenstate of $N_a(\mathbf{k})$ with eigenvalue $n_{\mathbf{k}} - 1$. However, if we require all multiparticle states to be normalized, so that $\langle n_{\mathbf{k}}|n_{\mathbf{k}}\rangle = 1$, the states $a^\dagger(\mathbf{k})|n_{\mathbf{k}}\rangle$ and $a(\mathbf{k})|n_{\mathbf{k}}\rangle$ do not produce normalized states. Rather, we have

$$(3) \quad a^\dagger(\mathbf{k})|n_{\mathbf{k}}\rangle = A|n_{\mathbf{k}} + 1\rangle$$

$$(4) \quad a(\mathbf{k})|n_{\mathbf{k}}\rangle = B|n_{\mathbf{k}} - 1\rangle$$

for some constants A and B that are determined by requiring normalization.

To find A and B , we can take the modulus of the states above. We get (we'll leave off the (\mathbf{k}) dependence of the a^\dagger and a operators to save typing; everything in what follows occurs at wave number \mathbf{k} ; we'll also assume A and B are real, although they could also be multiplied by some phase factor $e^{i\alpha}$, but this just complicates things unnecessarily). By using the commutation relation

$$(5) \quad [a, a^\dagger] = 1$$

we get, from 3

$$(6) \quad \langle n_{\mathbf{k}} | aa^\dagger | n_{\mathbf{k}} \rangle = A^2 \langle n_{\mathbf{k}} + 1 | n_{\mathbf{k}} + 1 \rangle = A^2$$

$$(7) \quad \langle n_{\mathbf{k}} | aa^\dagger | n_{\mathbf{k}} \rangle = \langle n_{\mathbf{k}} | a^\dagger a + 1 | n_{\mathbf{k}} \rangle$$

$$(8) \quad = \langle n_{\mathbf{k}} | N_a + 1 | n_{\mathbf{k}} \rangle$$

$$(9) \quad = (n_{\mathbf{k}} + 1) \langle n_{\mathbf{k}} | n_{\mathbf{k}} \rangle$$

$$(10) \quad = (n_{\mathbf{k}} + 1)$$

$$(11) \quad A = \sqrt{n_{\mathbf{k}} + 1}$$

Therefore

$$(12) \quad a^\dagger(\mathbf{k}) | n_{\mathbf{k}} \rangle = \sqrt{n_{\mathbf{k}} + 1} | n_{\mathbf{k}} + 1 \rangle$$

For the annihilation operator, we have from 4:

$$(13) \quad \langle n_{\mathbf{k}} | a^\dagger a | n_{\mathbf{k}} \rangle = B^2 \langle n_{\mathbf{k}} - 1 | n_{\mathbf{k}} - 1 \rangle = B^2$$

$$(14) \quad \langle n_{\mathbf{k}} | a^\dagger a | n_{\mathbf{k}} \rangle = \langle n_{\mathbf{k}} | N_a | n_{\mathbf{k}} \rangle$$

$$(15) \quad = n_{\mathbf{k}} \langle n_{\mathbf{k}} | n_{\mathbf{k}} \rangle$$

$$(16) \quad = n_{\mathbf{k}}$$

$$(17) \quad B = \sqrt{n_{\mathbf{k}}}$$

Therefore

$$(18) \quad a(\mathbf{k}) | n_{\mathbf{k}} \rangle = \sqrt{n_{\mathbf{k}}} | n_{\mathbf{k}} - 1 \rangle$$

This relation implies that applying $a(\mathbf{k})$ to a state that contains no particles with energy $\omega_{\mathbf{k}}$ (that is, where $n_{\mathbf{k}} = 0$) produces 0. In particular, if we apply $a(\mathbf{k})$ to the vacuum state, we end up with no state at all:

$$(19) \quad a(\mathbf{k}) | 0 \rangle = 0$$

Note that $|0\rangle$ and 0 aren't the same thing: $|0\rangle$ is a quantum state with no particles in it, while 0 is mathematically zero, that is, nothing. As an analogy, $|0\rangle$ is like having a bucket with nothing in it, while 0 corresponds to removing the bucket as well.

We can repeat exactly the same calculations for the antiparticle operators b^\dagger and b and get the results

$$(20) \quad b^\dagger(\mathbf{k}) |\bar{n}_{\mathbf{k}}\rangle = \sqrt{\bar{n}_{\mathbf{k}} + 1} |\bar{n}_{\mathbf{k}} + 1\rangle$$

$$(21) \quad b(\mathbf{k}) |\bar{n}_{\mathbf{k}}\rangle = \sqrt{\bar{n}_{\mathbf{k}}} |\bar{n}_{\mathbf{k}} - 1\rangle$$

PINGBACKS

Pingback: Feynman propagator for scalar fields