

DERIVATIVES OF THE DELTA FUNCTION

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

As the Dirac delta function is essentially an infinitely high spike at a single point, it may seem odd that its derivatives can be defined. The derivatives are defined using the delta function's integral property

$$(1) \quad \int_{-\infty}^{\infty} f(x) \delta(x) dx = f(0)$$

Consider the integral involving the n th derivative $\delta^{(n)}(x)$ and apply integration by parts:

$$(2) \quad \int_{-\infty}^{\infty} f(x) \delta^{(n)}(x) dx = f(x) \delta^{(n-1)}(x) \Big|_{-\infty}^{\infty} - \int_{-\infty}^{\infty} f'(x) \delta^{(n-1)}(x) dx$$

The integrated term is taken to be zero, since the delta function itself is constant (at zero) for all $x \neq 0$, so all its derivatives are zero except at $x = 0$. Therefore $\delta^{(n-1)}(x) = 0$ at the limits $-\infty$ and ∞ . We're therefore left with

$$(3) \quad \int_{-\infty}^{\infty} f(x) \delta^{(n)}(x) dx = - \int_{-\infty}^{\infty} f'(x) \delta^{(n-1)}(x) dx$$

Since this is true for all functions $f(x)$, the integrands must be equal, so we get

$$(4) \quad f(x) \delta^{(n)}(x) = -f'(x) \delta^{(n-1)}(x)$$

A common case is the first derivative, which satisfies

$$(5) \quad f(x) \delta'(x) = -f'(x) \delta(x)$$

If $f(x) = x$, we get the relation

$$(6) \quad x \delta'(x) = -\delta(x)$$

By iterating 4, we get

$$(7) \quad f(x) \delta^{(n)}(x) = (-1)^n \delta(x) f^{(n)}(x)$$

Example 1. Suppose $f(x) = 4x^2 - 1$. Then

$$(8) \quad \int_{-\infty}^{\infty} (4x^2 - 1) \delta'(x-3) dx = - \int_{-\infty}^{\infty} 8x \delta(x-3) dx$$

$$(9) \quad = -24$$

Example 2. With $f(x) = x^n$ we have, using 7

$$(10) \quad x^n \delta^{(n)}(x) = (-1)^n n! \delta(x)$$

Another use of the derivative of the delta function occurs frequently in quantum mechanics. In this case, we are faced with the integral

$$(11) \quad \int \delta'(x-x') f(x') dx'$$

where the prime in δ' refers to a derivative with respect to x , not x' . Thus the variable in the derivative is *not* the same as the variable being integrated over, unlike the preceding cases. In this case, since only x (and not x') is visible outside the integral, we can move the derivative outside the integral and get

$$(12) \quad \int \delta'(x-x') f(x') dx' = \frac{d}{dx} \int \delta(x-x') f(x') dx'$$

$$(13) \quad = f'(x)$$

Notice that in this case, there is no minus sign attached to the f' unlike in 5.

PINGBACKS

Pingback: [Differential operators - matrix elements and hermiticity](#)

Pingback: [Differential operator - eigenvalues and eigenstates](#)

Pingback: [Postulates of quantum mechanics: momentum](#)

Pingback: [Free particle in the position basis](#)