

HERMITIAN OPERATORS - COMMON EIGENFUNCTIONS IMPLIES THEY COMMUTE

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog and include the title or URL of this post in your comment.

Post date: 25 July 2021.

The eigenfunctions of a hermitian operator form a complete set, in the sense that any function can be expressed as a linear combination of these eigenfunctions. A consequence of this is the theorem that, if two hermitian operators have the *same* set of eigenfunctions (though possibly with different eigenvalues), then these two operators must commute.

If the two operators \hat{P} and \hat{Q} have the same complete set of common eigenfunctions, then a function f in Hilbert space can be written as a series in terms of these eigenfunctions

$$f = \sum c_n f_n \quad (1)$$

where f_n are the eigenfunctions, and c_n are the coefficients.

Now for operator \hat{P} we have $\hat{P}f_n = p_n f_n$ where p_n is the eigenvalue when \hat{P} operates on f_n . Similarly for \hat{Q} we have $\hat{Q}f_n = q_n f_n$. Since the two operators share the same set of eigenfunctions, applying the operators in either order to f will result in each term in the series being multiplied by the product of the two eigenvalues for each of the operators:

$$\hat{P}\hat{Q}f = \hat{Q}\hat{P}f = \sum c_n p_n q_n f_n \quad (2)$$

Thus the commutator will be zero if two operators share the same set of eigenfunctions. QED.