

LINEAR FUNCTIONALS AND ADJOINT OPERATORS

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog and include the title or URL of this post in your comment.

Post date: 3 Jan 2021.

1. LINEAR FUNCTIONALS

A *linear functional* is a linear map $\phi(v)$ from a vector space V to the number field \mathbb{F} which satisfies the two properties

- (1) $\phi(v_1 + v_2) = \phi(v_1) + \phi(v_2)$, with $v_1, v_2 \in V$.
- (2) $\phi(av) = a\phi(v)$ for all $v \in V$ and $a \in \mathbb{F}$.

That is, a linear functional acts on a vector and produces a number (either real or complex, depending on \mathbb{F}) as output.

A linear functional is actually a vector space, since it satisfies all the required axioms. Many of these axioms are satisfied because the V on which ϕ acts is a vector space. The only axiom that requires a bit of examination is the existence of an additive identity. This requires $\phi(v + 0) = \phi(v)$. From property 1 above, this means that $\phi(0) = 0$.

From the definition above, we can prove that any linear functional can be written as an inner product.

Theorem 1. *For any linear functional ϕ on V there is a unique vector $u \in V$ such that $\phi(v) = \langle u, v \rangle$ for all $v \in V$.*

Proof. We can write any vector v in terms of an orthonormal basis (e_1, \dots, e_n) as

$$v = \sum_{i=1}^n \langle e_i, v \rangle e_i \tag{1}$$

By applying the two properties of a linear functional above, we have

$$\phi(v) = \phi\left(\sum_{i=1}^n \langle e_i, v \rangle e_i\right) \tag{2}$$

$$= \sum_{i=1}^n \langle e_i, v \rangle \phi(e_i) \tag{3}$$

$$= \sum_{i=1}^n \langle e_i, \phi(e_i)v \rangle \tag{4}$$

$$= \sum_{i=1}^n \langle \phi(e_i)^* e_i, v \rangle \tag{5}$$

$$= \left\langle \left[\sum_{i=1}^n \phi(e_i)^* e_i \right], v \right\rangle \tag{6}$$

$$\equiv \langle u, v \rangle \tag{7}$$

where

$$u \equiv \sum_{i=1}^n \phi(e_i)^* e_i \tag{8}$$

We were able to move $\phi(e_i)$ inside the inner product in the third line above since $\phi(e_i)$ is just a number. To get the fourth line, we used

$$\langle \lambda^* u, v \rangle = \lambda \langle u, v \rangle = \langle u, \lambda v \rangle \tag{9}$$

which we have from our discussion of inner products.

To prove that u is unique, as usual we suppose there is another u' that gives the same result as u for all $v \in V$. This means that $\langle u', v \rangle = \langle u, v \rangle$ or $\langle u' - u, v \rangle = 0$ for all v . We can then choose $v = u' - u$, giving $\langle u' - u, u' - u \rangle = 0$, which implies that $u' - u = 0$ so $u' = u$. \square

2. ADJOINT OPERATORS

Suppose we have some linear operator T and some fixed vector u . We can then form the inner product

$$\phi(v) = \langle u, Tv \rangle \tag{10}$$

$\phi(v)$ is a linear functional since it satisfies the two properties specified earlier. It is now stated in Zwiebach's notes that because ϕ is a linear functional, we can write it in the form $\langle w, v \rangle$ for some vector w . It's not clear to me that this follows directly, since the original definition of a linear functional applied to the entire vector space V , whereas here we don't know whether T is a surjective operator, that is, whether the range of Tv is the

entire space V . The motivation behind this step is the definition of the adjoint operator, but in Axler's book (chapter 7.A), an adjoint is just defined directly without any motivation from linear functionals.

Anyway, we'll just go with Zwiebach's argument, since the rest of the derivation is fairly easy to follow. We assume that for a suitable vector w we have

$$\langle u, Tv \rangle = \langle w, v \rangle \quad (11)$$

The vector w depends on both the operator T and the vector u , so we can write it as a function of u , using the notation

$$w = T^\dagger u \quad (12)$$

This gives us the relation

$$\langle u, Tv \rangle = \langle T^\dagger u, v \rangle \quad (13)$$

At this stage, we can't be sure that T^\dagger is a linear operator; it may be some non-linear map from one vector to another. However, we have the following theorem.

Theorem 2. *The operator T^\dagger , called the adjoint of T , is a linear operator: $T^\dagger \in \mathcal{L}(V)$.*

Proof. Consider

$$\langle u_1 + u_2, Tv \rangle = \langle T^\dagger(u_1 + u_2), v \rangle \quad (14)$$

$$\langle u_1 + u_2, Tv \rangle = \langle u_1, Tv \rangle + \langle u_2, Tv \rangle \quad (15)$$

$$= \langle T^\dagger u_1, v \rangle + \langle T^\dagger u_2, v \rangle \quad (16)$$

$$= \langle T^\dagger u_1 + T^\dagger u_2, v \rangle \quad (17)$$

Comparing the first and last lines gives us

$$T^\dagger(u_1 + u_2) = T^\dagger u_1 + T^\dagger u_2 \quad (18)$$

A similar argument can be used for multiplication by a number:

$$\langle au, Tv \rangle = \langle T^\dagger(au), v \rangle \quad (19)$$

$$\langle au, Tv \rangle = a^* \langle u, Tv \rangle \quad (20)$$

$$= a^* \langle T^\dagger u, v \rangle \quad (21)$$

$$= \langle aT^\dagger u, v \rangle \quad (22)$$

Again, comparing the first and last lines we have

$$T^\dagger(au) = aT^\dagger u \quad (23)$$

Thus T^\dagger satisfies the two conditions required for linearity. \square

A couple of other results follow fairly easily (proofs are in Zwiebach's notes, if you're interested):

$$(ST)^\dagger = T^\dagger S^\dagger \quad (24)$$

$$(S^\dagger)^\dagger = S \quad (25)$$

A very important result is the representation of adjoint operators in matrix form.

If we have an orthonormal basis (e_1, \dots, e_n) and an operator T , then T transforms the basis according to (using the summation convention):

$$Te_k = T_{ik}e_i \quad (26)$$

Thus the matrix elements in this basis are found by taking the inner product with e_j :

$$\langle e_j, T_{ik}e_i \rangle = T_{ik} \langle e_j, e_i \rangle = T_{ik} \delta_{ji} = T_{jk} \quad (27)$$

Also,

$$T^\dagger e_j = T_{ij}^\dagger e_i \quad (28)$$

Taking the inner product on the right with e_k we get

$$\langle T^\dagger e_j, e_k \rangle = \langle T_{ij}^\dagger e_i, e_k \rangle = (T_{ij}^\dagger)^* \langle e_i, e_k \rangle = (T^\dagger)_{kj}^* \quad (29)$$

We can take the T_{ik} and T_{ik}^\dagger outside the inner product as they are just numbers. Using this, we have

$$\langle T^\dagger e_j, e_k \rangle = \langle e_j, T_{ik} e_i \rangle \quad (30)$$

$$\left(T^\dagger \right)_{kj}^* = T_{jk} \quad (31)$$

That is, in an orthonormal basis, the adjoint matrix is the complex conjugate transpose of the original matrix:

$$T^\dagger = (T^*)^T \quad (32)$$

The superscript T indicates 'transpose', not another operator!

REFERENCES

- (1) Axler, Sheldon (2015), *Linear Algebra Done Right*, 3rd edition, Springer. Chapters 3.F, 7.
- (2) Zwiebach, Barton, Online course *Mastering Quantum Mechanics Part 1: Wave Mechanics*. Archive available [here](#).
- (3) Liesen, Jörg & Mehrmann, Volker (2015), *Linear Algebra*, Chapter 13.
- (4) Shankar, R. (1994), *Principles of Quantum Mechanics*, Plenum Press, Chapter 1.

PINGBACKS

Pingback: Unitary operators