

GENERALIZED PRODUCT RULE: LEIBNIZ'S FORMULA

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.

We've seen that the product rule for derivatives is, for two functions $f(x)$ and $g(x)$:

$$(0.1) \quad (fg)' = f'g + fg'$$

This rule can be generalized to give Leibniz's formula for the n^{th} derivative of a product:

$$(0.2) \quad (fg)^{(n)} = \sum_{k=0}^n \binom{n}{k} f^{(k)} g^{(n-k)}$$

where the notation $f^{(n)} \equiv d^n f/dx^n$ and $\binom{n}{k} = \frac{n!}{k!(n-k)!}$ is the binomial coefficient.

The proof of this formula is a nice little exercise in proof by mathematical induction. An inductive proof requires two steps. First, we must show that the formula is true for one particular value of n , say $n = 0$. Second, we *assume* that the formula is true for a value $n = m$. From this assumption we then must prove that the formula is also true for $n = m + 1$. It doesn't matter what m is as long as it is taken to be equal to or greater than the value of n used in step 1. The idea is that if we can show that the truth of the formula for a particular value of n always implies its truth for the next value of n , then as long as we can demonstrate the truth of the formula for *some* value of n (as we do in step 1), the inductive reasoning implies it is true for all n greater than or equal to the specific value.

The first step in the proof is called the *anchor step* and the second step the *inductive step*.

In our case, taking $n = 0$ gives us the identity $(fg)^{(0)} = fg$ which is clearly true. So now we prove the inductive step. Assuming the formula is true for $n = m$ gives us

$$(0.3) \quad (fg)^{(m)} = \sum_{k=0}^m \binom{m}{k} f^{(k)} g^{(m-k)}$$

Taking the derivative of this, and applying the regular product rule, gives

$$(0.4) \quad (fg)^{(m+1)} = \sum_{k=0}^m \binom{m}{k} (f^{(k+1)}g^{(m-k)} + f^{(k)}g^{(m-k+1)})$$

In the first term, we can shift the summation index by replacing k by $k-1$ to get

$$(0.5) \quad (fg)^{(m+1)} = \sum_{k=1}^{m+1} \binom{m}{k-1} f^{(k)}g^{(m-k+1)} + \sum_{k=0}^m \binom{m}{k} f^{(k)}g^{(m-k+1)}$$

Note the limits on the two sums. We can now combine the sums in the regions where their indexes overlap to get

$$(0.6) \quad (fg)^{(m+1)} = fg^{(m+1)} + \sum_{k=1}^m \left[\binom{m}{k-1} + \binom{m}{k} \right] f^{(k)}g^{(m-k+1)} + f^{(m+1)}g$$

We can work out the sum of the two binomial coefficients by expanding them in terms of factorials:

$$(0.7) \quad \binom{m}{k-1} + \binom{m}{k} = \frac{m!}{(k-1)!(m-k+1)!} + \frac{m!}{k!(m-k)!}$$

Putting over a common denominator:

$$(0.8) \quad = \frac{(k)m!}{k!(m-k+1)!} + \frac{(m-k+1)m!}{k!(m-k+1)!}$$

$$(0.9) \quad = \frac{(m+1)!}{k!(m-k+1)!}$$

$$(0.10) \quad = \binom{m+1}{k}$$

Since $\binom{m+1}{0} = \binom{m+1}{m+1} = 1$ we can combine the two outlying terms in 0.6 to get the final result

$$(0.11) \quad (fg)^{(m+1)} = \sum_{k=0}^{m+1} \binom{m+1}{k} f^{(k)}g^{(m-k+1)}$$

This completes the inductive proof and establishes Leibniz's formula.

PINGBACKS

Pingback: [Associated Legendre functions](#)

Pingback: [Product rule and integration by parts](#)

Pingback: [Hermite polynomials – the Rodrigues formula](#)