

## LORENTZ TRANSFORMATIONS

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the auxiliary blog.


Although space-time diagrams can be used to analyze simple problems in relativity, a general algebraic transformation between two inertial frames is easier for most situations. These are the Lorentz transformations between the coordinates of two observers  $O_1$  and  $O_2$ , where  $O_2$  moves with speed  $v$  along the  $x_1$  axis of  $O_1$ .

We saw in an earlier post that coordinates perpendicular to the direction of motion are the same in both coordinate systems, so assuming that the transformation between the systems is linear (we show that this can be proved from more realistic assumptions in another post), the most general transformation is

$$\begin{aligned} (1) \quad & t_2 = at_1 + bx_1 \\ (2) \quad & x_2 = dt_1 + ex_1 \\ (3) \quad & y_2 = y_1 \\ (4) \quad & z_2 = z_1 \end{aligned}$$

where the coefficients  $a$ ,  $b$ ,  $d$  and  $e$  are functions of  $v$ .

We can use the following space-time diagram to work out these coefficients. In the diagram, the main axes are those of  $O_1$  with the axes of  $O_2$  drawn in green and red.



We've seen that the slope of the  $t_2$  axis is  $1/v$  and of the  $x_2$  axis is  $v$ . So when  $x_2 = 0$ , we must have for the  $t_2$  axis:

$$\begin{aligned}
 (5) \quad dt_1 + ex_1 &= 0 \\
 (6) \quad t_1 &= \frac{1}{v}x_1 \\
 (7) \quad &= -\frac{e}{d}x_1 \\
 (8) \quad -v &= \frac{d}{e}
 \end{aligned}$$

Similarly, when  $t_2 = 0$ , we must have for the  $x_2$  axis:

$$\begin{aligned}
 (9) \quad at_1 + bx_1 &= 0 \\
 (10) \quad t_1 &= vx_1 \\
 (11) \quad &= -\frac{b}{a}x_1 \\
 (12) \quad -v &= \frac{b}{a}
 \end{aligned}$$

We can now write the transformations in terms of two coefficients.

$$(13) \quad t_2 = a(t_1 - vx_1)$$

$$(14) \quad x_2 = e(x_1 - vt_1)$$

From the diagram, for the  $x_2$  axis, if we pick  $x_2 = 1$  we get from the second equation above:

$$(15) \quad 1 = e(\cos \alpha - v \sin \alpha)$$

Similarly, for the  $t_2$  axis, if we pick  $t_2 = 1$  we get from the first equation above

$$(16) \quad 1 = a(\cos \alpha - v \sin \alpha)$$

Comparing these two equations shows that

$$(17) \quad a = e$$

so we have

$$(18) \quad t_2 = a(t_1 - vx_1)$$

$$(19) \quad x_2 = a(x_1 - vt_1)$$

For a pair of events, one of which is the origin, the invariance of the interval can be used to say

$$(20) \quad x_2^2 - t_2^2 = x_1^2 - t_1^2$$

$$(21) \quad = a^2(x_1^2 - 2vx_1t_1 + v^2t_1^2 - t_1^2 + 2vt_1x_1 - v^2x_1^2)$$

$$(22) \quad = a^2(1 - v^2)(x_1^2 - t_1^2)$$

$$(23) \quad a = \pm \frac{1}{\sqrt{1 - v^2}}$$

We must choose the + sign so that at  $v = 0$  we get an identity.

This gives us the final form for the Lorentz transformations

$$(24) \quad t_2 = \frac{1}{\sqrt{1 - v^2}}(t_1 - vx_1)$$

$$(25) \quad x_2 = \frac{1}{\sqrt{1 - v^2}}(x_1 - vt_1)$$

$$(26) \quad y_2 = y_1$$

$$(27) \quad z_2 = z_1$$

These transformations apply to the special case of inertial frames that are aligned so that  $O_2$  travels with speed  $v$  along  $O_1$ 's  $x_1$  axis, with the corresponding  $y$  and  $z$  axes parallel in the two systems. This includes all real-life situations, although in some cases the coordinate systems may be

more complex and the transformations take on a more involved form. However, all the standard relativistic effects such as time dilation and length contraction can be derived from these Lorentz transformations.

#### PINGBACKS

[Pingback: Doppler effect](#)

[Pingback: Four-vectors: basics](#)

[Pingback: Four-velocity, momentum and energy](#)

[Pingback: Lorentz transformations as a linear map](#)

[Pingback: Four-velocity: an example](#)

[Pingback: Composition of velocities in relativity](#)