

FOUR-MOMENTUM CONSERVATION: A TRIP TO ALPHA CENTAURI

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

Reference: Moore, Thomas A., *A General Relativity Workbook*, University Science Books (2013) - Chapter 3; Problem 3.6.

As a rather fanciful example of using the conservation of four-momentum, suppose we have a spaceship of total mass (including fuel) M , initially at rest on Earth. The fuel consists of matter/anti-matter which when mixed, produces photons that are ejected out of the back of the ship. If the ship burns enough fuel to accelerate to $v = 0.95$, then travels to some star system such as Alpha Centauri, then decelerates to zero for a landing, then, after some time at its destination it reverses the trip by again accelerating to $v = 0.95$, returning to Earth, and decelerating to rest, what is its final mass as a fraction of its initial mass?

We can assume that all motion takes place along the x axis, and treat the problem in the Earth's frame. Then the initial momentum is

$$(0.1) \quad \mathbf{p}_0 = [M, 0]$$

After accelerating, the combined momentum of the ship + ejected photons is

$$(0.2) \quad \mathbf{p}_1 = \gamma m_1 [1, 0.95] + E_1 [1, -1]$$

where $\gamma = 1/\sqrt{1-v^2} = 3.2$, E_1 is the energy of the ejected photons and m_1 is the mass of the ship after burning the fuel needed to accelerate.

By the conservation of momentum, we have $\mathbf{p}_1 = \mathbf{p}_0$ so

$$(0.3) \quad \gamma m_1 + E_1 = M$$

$$(0.4) \quad 0.95\gamma m_1 - E_1 = 0$$

Adding these 2 equations, we get

$$(0.5) \quad m_1 = \frac{M}{1.95\gamma} = 0.16M$$

Now to decelerate the ship, we eject the photons ahead of the ship, and we start with a momentum of $\gamma m_1 [1, 0.95]$. After deceleration, the mass is now m_2 and the ship is at rest, so we must have

$$(0.6) \quad \mathbf{p}_2 = [m_2, 0] + E_2 [1, 1]$$

Again, conservation of momentum requires $\mathbf{p}_2 = \gamma m_1 [1, 0.95]$, so

$$(0.7) \quad \gamma m_1 = m_2 + E_2$$

$$(0.8) \quad 0.95\gamma m_1 = E_2$$

Subtracting these equations we get

$$(0.9) \quad m_2 = 0.05\gamma m_1$$

$$(0.10) \quad = \frac{0.05}{1.95} M$$

On the return trip, we go through exactly the same procedure, except we now start with a mass m_2 rather than M . Thus on the return to Earth, the ship's mass will be

$$(0.11) \quad m_E = \frac{0.05}{1.95} m_2$$

$$(0.12) \quad = \left(\frac{0.05}{1.95} \right)^2 M$$

Thus the ship's initial mass is

$$(0.13) \quad M = 1521 m_E$$

Virtually all the initial mass is fuel.

Incidentally, if we do this calculation for an arbitrary velocity v , we get

$$(0.14) \quad m_1 = \frac{M}{(1+v)\gamma}$$

$$(0.15) \quad = \sqrt{\frac{1-v}{1+v}} M$$

$$(0.16) \quad m_2 = \frac{1-v}{1+v} M$$

$$(0.17) \quad m_E = \left(\frac{1-v}{1+v} \right)^2 M$$

Thus each acceleration or deceleration multiplies the previous mass by a factor of $\sqrt{\frac{1-v}{1+v}}$.