

METRIC TENSOR: SINUSOIDAL COORDINATES

Link to: physicspages home page.

To leave a comment or report an error, please use the auxiliary blog.

Reference: Moore, Thomas A., *A General Relativity Workbook*, University Science Books (2013) - Chapter 5; Problem 5.5.

As another example of a different coordinate system, we can define a sinusoidal system by introducing coordinates u and w defined as:

$$\begin{aligned}(1) \quad & u = x \\(2) \quad & w = y - A \sin(bx)\end{aligned}$$

where A and b are constants. If x and y have units of length, then b must have units of length^{-1} . If A has units of length, then u and w both have units of length.

The curves of constant u are just the same as the curves of constant x , that is, vertical lines. The curves of constant w are defined by $y = k + A \sin(bx)$. These are parallel horizontal sine curves displaced vertically by the constant k .

The inverted relations are

$$\begin{aligned}(3) \quad & x = u \\(4) \quad & y = w + A \sin(bu)\end{aligned}$$

We can work out the metric of the sinusoidal system from the rectangular metric by direct calculation

$$(5) \quad g'_{ij} = g_{kl} \frac{\partial x^k}{\partial x'^i} \frac{\partial x^l}{\partial x'^j} = \begin{bmatrix} 1 + [Ab \cos(bu)]^2 & Ab \cos(bu) \\ Ab \cos(bu) & 1 \end{bmatrix}$$

Incidentally, we can check this by calculating the reverse transformation metric:

$$(6) \quad g_{ij} = g_{kl} \frac{\partial x'^k}{\partial x^i} \frac{\partial x'^l}{\partial x^j}$$

For example (using $x = u$):

(7)

$$g_{xx} = 1 + [Ab \cos(bx)]^2 - [Ab \cos(bx)]^2 - [Ab \cos(bx)]^2 + [Ab \cos(bx)]^2$$

(8) = 1

Similar calculations give the other components of the rectangular metric.

Now suppose we have an object moving with velocity \mathbf{v} such that $v^x = v$ and $v^y = 0$. In the sinusoidal system, we get

$$(9) \quad v^u = \frac{\partial u}{\partial x^i} v^i$$

$$(10) \quad = v$$

$$(11) \quad v^w = \frac{\partial w}{\partial x^i} v^i$$

$$(12) \quad = -Ab \cos(bu) v$$

$$(13) \quad = -Ab \cos(bvt) v$$

where the last line arises because $u = x = vt$, where t is the time.

The square is

(14)

$$v^2 = g'_{ij} v^i v^j$$

(15)

$$= \left(1 + [Ab \cos(bu)]^2\right) v^2 - [Ab \cos(bu)]^2 v^2 - [Ab \cos(bu)]^2 v^2 + [Ab \cos(bu)]^2 v^2$$

(16)

$$= v^2$$

Thus the square of the velocity is invariant.

The unit vector \mathbf{e}_u is tangent to the constant- w curves and points in the direction of increasing u . Since the constant- w curves are sine curves, this unit vector starts off horizontal (when $bu = \pi/2$), then slopes downward as bu heads towards $3\pi/2$ at which point it is horizontal again. Then it slopes upwards as bu heads towards $5\pi/2$ and so on. Its magnitude is given by

$$|\mathbf{e}_u| = \sqrt{g_{uu}} = \sqrt{1 + [Ab \cos(bu)]^2}.$$

The other unit vector \mathbf{e}_w is tangent to the constant- u curves, so it always points up, and always has a length of 1. Note that the off-diagonal elements of g'_{ij} are zero when bu is an odd multiple of $\pi/2$, which is where \mathbf{e}_u is horizontal, and thus perpendicular to \mathbf{e}_w , so $g'_{uw} = g'_{wu} = \mathbf{e}_u \cdot \mathbf{e}_w = 0$.

The reason that v^w is not constant, even though the velocity itself is constant is because the unit vector \mathbf{e}_u oscillates in both magnitude and direction. Since \mathbf{e}_w is a constant, the component multiplying this unit vector (that is, v^w) must also oscillate to compensate for the non-constant u component.

In the rectangular system, the acceleration is zero (since v is constant). In the sinusoidal system, $dv^u/dt = 0$, but $dv^w/dt \neq 0$. Thus the magnitude of a^2 would not be invariant under the transformation of coordinates, so this cannot be the correct way of calculating derivatives in a general coordinate system.

PINGBACKS

Pingback: [Metric tensor: spherical coordinates](#)

Pingback: [Christoffel symbols in sinusoidal coordinates](#)