

GEODESIC EQUATION: 2-D SPACE-TIME

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

Reference: Moore, Thomas A., *A General Relativity Workbook*, University Science Books (2013) - Chapter 8; Problem 8.5.

This is an example of the geodesic equation in a 2-d space-time (with one time and one space dimension). The metric is given in a general way as

$$ds^2 = -dt^2 + f^2(q) dq^2 \quad (1)$$

where q is the generalized spatial coordinate, and f is an arbitrary function. The metric tensor is then

$$g_{ij} = \begin{bmatrix} -1 & 0 \\ 0 & f^2(q) \end{bmatrix} \quad (2)$$

Using the time component of the geodesic equation, we set $a = t$ in:

$$\frac{d}{d\tau} \left(g_{aj} \frac{dx^j}{d\tau} \right) - \frac{1}{2} \partial_a g_{ij} \frac{dx^i}{d\tau} \frac{dx^j}{d\tau} = 0 \quad (3)$$

We get

$$-\frac{d^2t}{d\tau^2} = 0 \quad (4)$$

From this we conclude that

$$\frac{dt}{d\tau} = k \quad (5)$$

for some constant k .

Using the condition $\mathbf{u} \cdot \mathbf{u} = -1$ we get

$$g_{ij} \frac{dx^i}{d\tau} \frac{dx^j}{d\tau} = - \left(\frac{dt}{d\tau} \right)^2 + f^2 \left(\frac{dq}{d\tau} \right)^2 \quad (6)$$

$$= -k^2 + f^2 \left(\frac{dq}{d\tau} \right)^2 \quad (7)$$

$$= -1 \quad (8)$$

$$\frac{dq}{d\tau} = \pm \frac{1}{f} \sqrt{(k^2 - 1)} \quad (9)$$

We can write this in terms of dq/dt :

$$\frac{dq}{d\tau} = \frac{dq}{dt} \frac{dt}{d\tau} \quad (10)$$

$$= k \frac{dq}{dt} \quad (11)$$

$$\frac{dq}{dt} = \pm \frac{1}{f(q)} \sqrt{\left(1 - \frac{1}{k^2}\right)} \quad (12)$$

That is, the geodesic is the solution of this differential equation.

If we define a new coordinate system in which $t' = t$ and $q' \equiv F(q)$ is the antiderivative (integral) of $f(q)$ then we can transform the metric tensor to this new coordinate system using the standard transformation formula

$$g'_{ij} = \frac{\partial x^k}{\partial x'^i} \frac{\partial x^l}{\partial x'^j} g_{kl} \quad (13)$$

By implicit differentiation:

$$\frac{\partial q'}{\partial q'} = \frac{\partial F}{\partial q} \frac{\partial q}{\partial q'} + \frac{\partial F}{\partial t} \frac{\partial t}{\partial q'} \quad (14)$$

$$1 = f(q) \frac{\partial q}{\partial q'} + 0 \quad (15)$$

$$\frac{\partial q}{\partial q'} = \frac{1}{f(q)} \quad (16)$$

The only other non-zero derivative is

$$\frac{\partial t}{\partial t'} = 1 \quad (17)$$

The new metric is therefore

$$g'_{ij} = \begin{bmatrix} -1 \left(\frac{\partial t}{\partial t'}\right)^2 & 0 \\ 0 & f^2(q) \left(\frac{\partial q}{\partial q'}\right)^2 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \quad (18)$$

This is the metric of flat space-time in rectangular coordinates. Thus any metric with $g_{tt} = -1$ represents flat space-time, since $f(q)$ is just a transformation of the flat metric using different coordinates.