

PARTICLE FALLING TOWARDS A MASS: TWO TYPES OF VELOCITY

Link to: [physicspages home page](#).

To leave a comment or report an error, please use the [auxiliary blog](#).

Reference: Moore, Thomas A., *A General Relativity Workbook*, University Science Books (2013) - Chapter 10; Problem 10.4.

The radial equation of motion in the Schwarzschild metric is

$$(0.1) \quad \frac{1}{2} \left(\frac{dr}{d\tau} \right)^2 + \frac{1}{2} \frac{l^2}{r^2} - GM \left(\frac{1}{r} + \frac{l^2}{r^3} \right) = \frac{1}{2} (e^2 - 1)$$

We can use this to derive an equation for dr/dt , the rate of change of r with respect to the Schwarzschild time coordinate. The coordinate t isn't the time as measured by any particular object (that time is the proper time τ in the reference frame of the object) so we wouldn't expect it to be the same as $dr/d\tau$.

To get the equation, we can use

$$(0.2) \quad \frac{dr}{d\tau} = \frac{dr}{dt} \frac{dt}{d\tau}$$

$$(0.3) \quad = \frac{dr}{dt} e \left(1 - \frac{2GM}{r} \right)^{-1}$$

where the last line uses the definition of e . Plugging this into the top equation we get

$$(0.4) \quad \frac{dr}{dt} = \frac{1}{e} \left(1 - \frac{2GM}{r} \right) \left[e^2 - 1 + 2GM \left(\frac{1}{r} + \frac{l^2}{r^3} \right) - \frac{l^2}{r^2} \right]^{1/2}$$

As r approaches $2GM$, $dr/dt \rightarrow 0$.

In the special case where we drop an object from rest at $r = r_0$, we can work out both dr/dt and $dr/d\tau$. In this case, motion is radially inward so $l = 0$. To find e , we use the fact that for an object at rest at $r = r_0$:

$$(0.5) \quad e = \left(1 - \frac{2GM}{r_0}\right) \frac{dt}{d\tau}$$

$$(0.6) \quad = \left(1 - \frac{2GM}{r_0}\right) u^t$$

$$(0.7) \quad = \left(1 - \frac{2GM}{r_0}\right) \left(1 - \frac{2GM}{r_0}\right)^{-1/2}$$

$$(0.8) \quad = \left(1 - \frac{2GM}{r_0}\right)^{1/2}$$

We have therefore

$$(0.9) \quad \frac{dr}{dt} = \frac{1}{e} \left(1 - \frac{2GM}{r}\right) \left(e^2 - 1 + \frac{2GM}{r}\right)^{1/2}$$

$$(0.10) \quad = \frac{1 - \frac{2GM}{r}}{\sqrt{1 - \frac{2GM}{r_0}}} \sqrt{2GM \left(\frac{1}{r} - \frac{1}{r_0}\right)}$$

$$(0.11) \quad = \left(1 - \frac{2GM}{r}\right) \sqrt{\frac{2GM}{r}} \sqrt{\frac{r_0 - r}{r_0 - 2GM}}$$

From 0.1 we get

$$(0.12) \quad \frac{dr}{d\tau} = \sqrt{e^2 - 1 + \frac{2GM}{r}}$$

$$(0.13) \quad = \sqrt{2GM \left(\frac{1}{r} - \frac{1}{r_0}\right)}$$

Comparing the two, we see that

$$(0.14) \quad \frac{dr}{dt} = \frac{1 - \frac{2GM}{r}}{\sqrt{1 - \frac{2GM}{r_0}}} \frac{dr}{d\tau}$$

For the case where the object is released from rest at $r_0 = \infty$, the speed at $r = 6GM$ is

$$(0.15) \quad \frac{dr}{d\tau} = \frac{1}{\sqrt{3}}$$

which agrees with the earlier calculation done by a different method.